

## گزارش و مترادف جدید در جنس *Ranunculus* برای ایران

### A new record and a synonym in the genus *Ranunculus* (Ranunculaceae) from Iran

Received: 01.05.2010 / Accepted: 16.06.2010

دریافت: ۱۳۸۹/۲/۱۱ / پذیرش: ۱۳۸۹/۳/۲۶

**M. Pakravan✉:** Assistant Prof., Department of Biology,  
Faculty of Science, Alzahra University, Tehran, Iran  
(E-mail: mpf176@yahoo.com)

In a project on the Flora of Iran (Trib.: Ranunculeae), many herbarium specimens of the genus *Ranunculus* were studied. The results of this study are a new record and suggestion of a synonym for the Flora of Iran, specimens of which are preserved in the herbarium of Research Institute of Forest & Rangelands (TARI).

#### ***Ranunculus meyerianus* Rupr.**

This species was first described from Crimea (Rupr., Fl. Cauc. I, 1869). It is distributed from Europe through Caucasia to N. Iran. During the studies of *Ranunculus* species collected from Arasbaran protected region, one specimen of Kharil mountains collected by Assadi & Maasoumi (TARI 20266) was seen with the following characters:

Perennial, rootstock short, indumentums of basal leaves' petioles similar to the stems, composed of spreading hairs, blade tripartite to base, the middle segment subsessile, more or less broadly rhombic, tripartite to the middle or beyond, with connate incised lobes, or with dentate incised segments, covered with long appressed hairs on both surfaces especially lower surface. Flowers large, sepals oval-elliptic, densely covered with spreading yellow hairs. Achens 2.5-3.5 mm long, oval-oblong, smooth with narrow border, laterally compressed, passing gradually into a short proximally broad beak, 0.2-0.5 mm long, triangle in outline, straight.

This species is closely related to *R. polyanthemos* L. by almost straight broad beak of its achen (0.5-1 mm long, uncinate in *R. polyanthemos*) and by denser and longer hairs on the stems, petioles and leaves, also difference in ± sessile middle segment of leaf (stalked in *R. polyanthemos*). It also resembles *R. grandiflorus* L., but differentiated in having narrower leaf segments, broad short and straight beak (in *R. meyerianus*) and spread sepals (reflexed in *R. grandiflorus*). Parsa & Maleki (1986) have reported this species from Lorestan (W Iran) without mentioning any herbarium number. Also Assadi (1998) has recorded this species from Arasbaran protected region but it was misidentified. These specimens are recognized as follows: 20974, 24361 as *R. grandiflorus* and 20403, 24752 as *R. polyanthemos*. The achens shapes are shown in Fig. 1 and the characters are compared in Table 1.

#### New synonym

##### ***R. constantinopolitanus* (DC.) d'Urv.**

Syn.: *R. farsicus* Rech. F.; *R. constantinopolitanus* (DC.) D'Urv. f.

Rechinger (1951) recognized *R. farsicus* from *R. constantinopolitanus* by having narrow achen and uncinate beak, short stem and narrow leaf segment. However, through author's detailed field examination and measurement of several herbarium materials together with studying photo type, many intermediate forms were found that the distinction between them was not sharp enough. Moreover, *R. constantinopolitanus* shows many polymorphisms in size, shape of achen beak and leaves.

**منیژه پاکروان✉:** استادیار دانشکده علوم پایه، دانشگاه الزهراء (س)، تهران  
(E-mail: mpf176@yahoo.com)


طی بررسی های چهار ساله روی طرح فلور ایران (طایفه Ranunculeae)، نمونه های متعددی از هرباریوم های مختلف بررسی شد. این مطالعه منتج به گزارش یک گونه و مترادف کردن یک گونه شد که نمونه های آن در هرباریوم مؤسسه تحقیقات جنگل ها و مراتع ("TARI") موجود است.

##### ***Ranunculus meyerianus* Rupr.**

این گونه اولین بار در فلور قفقاز از کریمه شرح داده شده است (Rupr., 1869, Fl. Cauc. I) و از اروپا تا قفقاز و شمال ایران گسترش یافته است. طی بررسی نمونه های جمع آوری شده از جنس آله (Ranunculus) از منطقه حفاظت شده ارسباران، نمونه ای از کوه خریل، ۲۰۰۰-۲۵۰۰ متر، اسدی و معصومی (TARI ۲۰۲۶۶)، مشاهده شد که دارای مشخصات زیر بود:

گیاهی چندساله با ریشه کوتاه، برگ های قاعده ای با دمبرگی بلند که مانند ساقه پوشیده از کرک های گستردگی است. پهنک برگ سه بخشی تا قاعده، بخش میانی تقریباً بدون دمبرگ، کم و بیش لوزی پهن، سه بخشی تا میانه یا بیشتر از میانه، با بریدگی یا با قطعات دندانه دار، پوشیده از کرک های بلند و خوابیده در دو سطح به خصوص در سطح زیرین. گل ها بزرگ، زرد رنگ. کاسبرگ ها تخم مرغی- بیضوی با پوششی مترادکم از کرک های گستردگی. فندقه به طول ۲/۵-۳/۵ میلی متر، تخم مرغی- مستطیلی، بدون کرک با حاشیه باریک، در جوانب دارای فشردگی، منتهی به یک منقار کوتاه، راست و مثلثی و پهن به طول ۰/۵-۰/۲ میلی متر.

این گونه به *R. polyanthemos* L. نزدیک است و با داشتن منقار پهن و راست در میوه (طول منقار در ۰/۵-۱ میلی متر است) و کرک های مترادکم و بلند روی ساقه، دمبرگ و پهنک و نیز با قطعه میانی کم و بیش بدون پایه در برگ (در *R. polyanthemos* پایه دار است) از آن متمایز می شود. همچنین گونه فوق شباهت هایی به *R. grandiflorus* دارد اما با داشتن برگ هایی با قطعات باریک تر، منقار کوتاه، پهن و راست و کاسبرگ های گستردگی (در *R. grandiflorus* دارای برگ شتگی به بیرون است) مجزا می شود. پارسا و ملکی (۱۹۸۶) این گونه را از لرستان ذکر کرده اند ولی هیچ شماره هرباریومی و نام هرباریوم برای آن بیان نکرده اند. همچنین اسدی (۱۹۹۸) این گونه را از منطقه حفاظت شده ارسباران ذکر کرده است ولی نمونه ها درست شناسایی نشده اند. این نمونه ها مجدداً به شرح زیر شناسایی شدند: نمونه های *R. grandiflorus* و نمونه های ۲۴۳۶۱ و ۲۰۹۷۴ به عنوان *R. grandiflorus* و نمونه های ۲۴۷۵۲ و ۲۰۴۰۳ به عنوان *R. polyanthemos* (شکل ۱ و جدول ۱).



شکل ۱- شکل‌های فندقه در گونه‌های *R. polyanthemos* A، *R. grandiflorus* C و B، (۲۰۴۰۳) *R. polyanthemos* .C و B، (۲۰۴۰۳) *R. meyerianus* F، (۲۴۷۵۲) *R. polyanthemos* E، (۲۱۶۶۴) *R. grandiflorus* D، (۲۴۳۶۱) *R. meyerianus* F، (۲۴۷۵۲) *R. polyanthemos* E، (۲۱۶۶۴) *R. grandiflorus* D، (۲۴۳۶۱)

Fig.1. Photographs of achenes in *Ranunculus* species: A. *R. polyanthemos* (20403), B & C. *R. grandiflorus* (24361), D. *R. grandiflorus* (21664), E. *R. polyanthemos* (24752), F. *R. meyerianus*.

جدول ۱- صفات ریخت‌شناسی در سه گونه *Ranunculus* spp.Table 1. Morphological characters of *Ranunculus* spp.

Species	Stem indumentum	Leaf indumentum	Leaf shape	Achen size (mm)	Beak shape	Beak length (mm)
<i>R. meyerianus</i> Rupr.	Dense, spreading, long hair	Long retrose hair	Tripartite, segments more divided into narrow lobes	2.5- 3.5×1.8- 2.3	Broad triangle straight	0.2-0.5
<i>R. polyanthemos</i> L.	White spreading hair	Appressed long hair	3-5 partite to base, segments deeply dissected	2.5-4×2-3	Slightly incurved	0.3-1
<i>R. grandiflorus</i> L.	Dense spreading hair	Dense appressed hair	Tripartite, segments 2-3 part	2.3- 2.7×1.8- 2.3	Incurved	0.3-0.7
<i>R. constantinopolitanus</i> (DC.) Urv.	Long spreading hirsute hair	Dense long appressed villose	Tripartite to beyond middle, segments broad triangle	1.5- 3.5×2.5-5	Hamate- involute	0.8-3.5
<i>R. farsicus</i> Rech. f.	Dense hirsute	Dense long appressed hair	Tripartite, segments narrow ±acute	3×2	Hamate	1

It has several polyploidy levels in chromosomes ( $2n=14, 42$ ) (Gregory 1941, Nemati 2007). But still more specimens with different size and shape are to be expected. *R. constantinopolitanus* has a widespread distribution from Europe to Caucasus, Syria, Iraq, North of Iran to central and south of Iran. The type locality of *R. farsicus* is Fars Province (Dashte-Arjan). The differences in size of leaf segments and shape are also misleading, as is found in several intermediate plants. Almost closed to the type locality, there are plants with taller stem and broader leaf segments (Esfahan: Semiroom, Vardasht-Garmab, 2350-2600 m, Nowroozi 3822; Kohkiloyeh & Boyerahmad: 27 km Sisakht, 1650, Jamzad *et al.* 69438). The beak of achen is 1 mm in *R. farsicus* and 0.7-2.5 mm in *R. constantinopolitanus*. It was, therefore, concluded that, *R. farsicus* is not a valid name and must be considered as a synonym of *R. constantinopolitanus*.

**Acknowledgements:** The author wish to thank Dr. M. Assadi for his valuable comments. I also wish to thank Mrs. F. Aghabeigi, former curator of IRAN herbarium for providing the herbarium specimens.

## References

- Assadi, M. 1998. Plants of Arasbaran protected area, NW Iran. *Iran. J. Bot.* 4(1): 1-59.  
 Gregory, W. 1941. Phylogenetic and cytological studies in the Ranunculaceae. *Trans. Amer. Philos. Soc.* 31.  
 Nemati, S. 2007. Systematic study of the genus *Ranunculus* (Group Praemorsa) in Iran. Unpublished MSc. Thesis; Alzahra University, Tehran, Iran.  
 Parsa, A. & Maleki, Z. 1986. Flora of Iran. Vol. 2: 166.  
 Rechinger, K.H. 1951. *Anz. Math.-Nat. Kl. Österr. Akad. Wiss.* 88: 222.

## مترادف جدید

***Ranunculus constantinopolitanus* (DC.) d'Urv.**  
 Syn.: *R. farsicus* Rech. f.; *R. constantinopolitanus* (DC.) D'Urv. f. *farsicus* (Rech. f.) Parsa  
 رشینگر (Rechinger 1951) گونه *R. farsicus* را با صفاتی مانند فندههای باریک با منقار داسی شکل، ساقه کوتاه و برگهایی با قطعات باریک از *R. constantinopolitanus* مجزا کرده است. طی مطالعات وسیعی که روی نمونههای گیاهی در محیط طبیعی و نیز روی نمونههای هرباریومی صورت گرفته، نمونههای بسیاری یافته گردید که دارای صفات حد واسطه بودند، به طوری که تشخیص آنها آن طور که رشینگر ذکر کرده است امکان پذیر نبود. به علاوه، *R. constantinopolitanus* دارای پلیمورفیسم زیادی در اندازه و شکل منقار میوه و برگ است. همچنین سطوح پلیپلوبیدی مختلفی در کروموزومهای ( $2n=14, 42$ ) آنها ذکر شده است (عمتی ۱۳۸۶ و ۱۹۴۱). بنابراین، انتظار می‌رود نمونههای بسیاری با اندازه‌ها و شکل‌های مختلفی در اندامهای ذکر شده یافته شود. از طرفی، *R. constantinopolitanus* پراکنش وسیعی داشته از اروپا تا قفقاز، سوریه، عراق، شمال، جنوب و مرکز ایران گسترش یافته است. محل نمونه تیپ *R. farsicus* استان فارس (دشت ارزن) است (تصویر نمونه تیپ برسی شد). تفاوت در اندازه قطعات برگ و شکل آنها صفاتی گمراه‌کننده است، زیرا گیاهان متعددی با صفات حد واسطه یافته می‌شوند. تقریباً نزدیک محل تیپ گونه مذکور، گیاهانی با ساقه بلندتر و برگهایی با قطعات پهن‌تر مشاهده کردید (اصفهان: سمیرم، وردشت-گرماب، ۲۳۵۰-۲۶۰۰ متر، سوروزی ۳۸۲۲ TARI) که گیلوبیه و بویراحمد: (TARI ۶۹۴۳۸) ۲۷ کیلومتری سی سخت، ۱۶۵۰ متر، جمزاد و همکاران (TARI ۶۹۴۳۸) طول منقار فنده در *R. farsicus* در حدود یک میلی‌متر ذکر شده است اما در *R. constantinopolitanus* از ۲/۵-۰ میلی‌متر تغییر می‌کند. بنابراین، *R. farsicus* نمی‌تواند گونه مجزایی باشد، لذا در اینجا به عنوان مترادف *R. constantinopolitanus* مورد ملاحظه قرار گرفته است. سپاسگزاری: از آقای دکتر مصطفی اسدی به خاطر راهنمایی‌های ارزندeshan در این مقاله سپاسگزاری می‌شود. همچنین از خانم مهندس فاطمه آقایی‌گی مسئول وقت هرباریوم "IRAN" برای در اختیار گذاشتن نمونههای هرباریومی تشکر می‌شود.