

Diversity and distribution patterns of *Solanaceae* in Iran: Implications for conservation and habitat management with emphasis on endemism and diversity in SW Asia

Received: 20.08.2016 / Accepted: 11.12.2016

Sadaf Sayadi: MSc Student, Department of Plant Sciences and Technology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, GC, Tehran, Iran

Ahmadreza Mehrabian✉: Assistant Prof., Department of Plant Sciences and Technology, Faculty of Life Sciences and Biotechnology, Shahid Beheshti University, GC, Tehran, Iran (a_mehrabian@sbu.ac.ir)

Abstract

The increasing rate of irreversible damages to natural habitats has reinforced the need for gathering biodiversity fundamental data to conservation. Priority setting is the most important step in protection programs. *Solanaceae* as the one of important economic families of eudicots distributed throughout North and South America, Europe, Africa, Asia, Australia, and the South Pacific. Up to now, little attention has been paid to the patterns and diversity centers of *Solanaceae* in Iran. The present study describes in as much detail as possible the spatial distribution, areas with greater species richness, and centers of diversity for taxa. An ecogeographical database was developed using the records that corresponds to 3123 herbarium specimens belong to 2560 localities. The localities were marked using ArcView version 3.2 (ESRI 2000) on geo-referenced maps (1/106) of Iran. The distribution patterns of the taxa were mapped per $1^\circ \times 1^\circ$ universal transverse Mercator grid cells (100 km² with the exception of boundary area). Threatened categories of Iranian *Solanaceae*, including CR (43.4%) and VU (34%), mainly (58.9%) distributed in Irano-Turanian phytocorion. *Solanaceae* represents the highest richness in the central Alborz, Zagros, Kopet Dagh and Makran Mountains based on recent criteria. The conservation value of habitats of *Solanaceae* in Iran ranges from 0.15 (in coastline of Persian Gulf) to 11.19 (in eastern Alborz).

Keywords: Conservation management, plant diversity, priorities for conservation, red list of species

الگوهای پراکنش تیره سیب‌زمینی در ایران: کاربرد در حفاظت و مدیریت رویشگاه‌ها

با تاکید بر اندمیسم و تنوع در جنوب غرب آسیا*

دریافت: ۱۳۹۵/۰۵/۳۰ / پذیرش: ۱۳۹۵/۰۹/۲۱

صدف صیادی: دانشجوی کارشناسی ارشد، گروه علوم و فناوری گیاهی، دانشکده علوم و فناوری زیستی، دانشگاه شهید بهشتی، تهران، ایران

احمدرضا محرابیان✉: استادیار گروه علوم و فناوری گیاهی، دانشکده علوم و فناوری زیستی، دانشگاه شهید بهشتی، تهران، ایران (a_mehrabian@sbu.ac.ir)

خلاصه

افزایش نرخ آسیب‌ها به زیستگاه‌های طبیعی نیاز به تدوین اطلاعات پایه جهت حفاظت را تقویت نموده است. الویت، مهمترین فاکتور در برنامه‌های حفاظتی می‌باشد. تیره سیب‌زمینی به عنوان یکی از گروه‌های دولپه‌ای حقیقی دارای طیف گسترده‌ای از مصارف دارویی، وابستگان وحشی گیاهان زراعی و زینتی می‌باشد. این تیره شامل ۵۵ گونه متعلق به نه جنس از شش طایفه در ایران است. مطالعه حاضر با هدف توصیف ویژگی‌های ژئوبوتانیکی، پراکنش فضایی، مناطق غنای گونه‌ای، مراکز تنوع و اندمیسم و همچنین گونه‌های نادر و در خطر انقراض با رویکرد ژئوبوتانیکی و حفاظت تبیین شده است. این پژوهش براساس ارزیابی ۳۱۲۳ نمونه ثبت شده از ۲۶۵۰ موقعیت جغرافیایی صورت پذیرفته است. این مناطق جغرافیایی در سلول‌های شبکه‌ای $1^\circ \times 1^\circ$ (UTM) با مقیاس ۱۰۰ کیلومتر مربع به استثنای منطقه مرزی برای تعیین الگوی پراکنش و مناطق مهم به منظور مدیریت حفاظت جهت تهیه نقشه‌ها مورد ارزیابی قرار داده شده است. طبقات تهدید آرایه‌های تیره سیب‌زمینی در ایران عبارتند از: بحرانی (۴۳/۴ درصد) و آسیب‌پذیر (۳۴ درصد) که عمده آن‌ها (۵۸/۹ درصد) در منطقه ایران-تورانی استقرار یافته است. نتایج این مطالعه نشان می‌دهد که بالاترین غنای گونه‌ای این تیره در کوه‌های البرز، زاگرس، کپه‌داغ و مکران تمرکز دارد. علاوه بر این، دامنه ارزش حفاظتی رویشگاه‌ها بین ۰/۱۵ (نوار ساحلی فارسی خلیج فارس و دریای عمان) تا ۱۱/۱۹ (البرز شرقی) متغیر می‌باشد.

واژه‌های کلیدی: الویت‌های حفاظتی، تنوع گیاهی، فهرست سرخ گونه‌ها، مدیریت حفاظت

* بخشی از پایان‌نامه کارشناسی نگارنده اول به راهنمایی دکتر احمدرضا محرابیان ارائه شده به دانشکده علوم و فناوری‌های زیستی دانشگاه شهید بهشتی، تهران

Introduction

The increasing rate of irreversible damage to natural habitats has reinforced the need for gathering biodiversity fundamental data to conservation planning (Mutke & Barthlott 2005). Setting priorities is the most important step in protection programs (Lovett *et al.* 2000). Up to now, many conservationists have focused on mapping the centers of endemism (Linder 2001, Jetz *et al.* 2004, Mittermeier *et al.* 2005), biodiversity hotspots (Williams *et al.* 1997, Reid 1998, Meyers *et al.* 2000), diversity centers of rare and endangered taxa (Dobson *et al.* 1997, Prendergast *et al.* 1993, Mittermeier *et al.* 1999), diversity centers of crop wild relatives (Iriando *et al.* 2008) as well as medicinal plants (Kala 2005, Robenson 2008).

In the conservation approach, all taxa are not of equal significance (Vane-Wright *et al.* 1991, Arponen *et al.* 2005). The vulnerability of a wide range of taxa makes it necessary to assess the important taxa groups (Meyers *et al.* 2000, Negi & Gadgil 2002). Taxon-based diversity mapping (Müller *et al.* 2003) has been used for plant taxa on regional (Wagnetiz 1986, Rechinger 1986, Bischler & Jovet-Ast 1996, Zaharieva *et al.* 2005, Luna Vega *et al.* 2004, Zhongqiang *et al.* 2006, Moreira & Munoz-Schick 2007, Dowe 2010, Marshal *et al.* 2012, Mehrabian 2015) and continental-global (Barthlott *et al.* 1996, Farjon 1998, Jaramillo & Manos 2001, Ortega Baes & Godinez-Alvarez 2006, Olmstead 2013, Laurence *et al.* 2013) scales. Herbarium accessions usually provide the best data accessible for target taxa (Schmidt *et al.* 2005, Applequist *et al.* 2007, Lavoie 2013). The data shows efficiency in distribution patterns and ecological modeling (Heyligers 1998, Crawford & Schulman *et al.* 2007, Hoagland 2009, Fuentes *et al.* 2010, Hiopkins 2007, Raes *et al.* 2009, Droisart *et al.* 2012). Besides, the geodiversity and diversity of topography, climate, and soils correlate with the diversity patterns of vascular plants (Barthlott *et al.* 1996, 2000, Faith & Walker 1996, Jedicke 2001, Braun *et al.* 2002).

Solanaceae as the one of important economic families of eudicots distributed throughout North and South America, Europe, Africa, Asia, Australia, and the

South Pacific (Ladiges *et al.* 2011). It comprise 90 genera with more than 3500 species (D'Arcy 1986) and include some species with medicinal usage and alkaloids resistant to herbivorous (Heiser 1969) and several crop wild relatives (Samuels 2015) that are the focus of pharmacologists, ecologists, and gene breeders. Some species of this family can be used ecologically to maintain the healthy performance of the ecosystem as surrogates for conservation (Heywood & Dullo 1996, Caro 2010). *Solanaceae* exhibits diverse life forms that include trees, shrubs, herbs, climbers, and epiphytes. Worldwide, some taxa are confined to specific ecological niches (D'Arcy 1986) for which it is necessary to provide distribution patterns at the regional scale to determine the centers of diversity, areas of endemism, and categorize their conservation status.

Up to now, little attention has been paid to the patterns and diversity centers of *Solanaceae* in Iran. The present study describes in as much detail as possible the spatial distribution, areas with greater species richness, centers of diversity for medicinal taxa and crop wild relatives, and rare and endangered taxa in a geobotanical and conservation approach. This study summarizes the results of a large-scale survey of diversity patterns of Iranian families in what is recognized as one of the most important diversity centers in the world.

- Study area

Iran covers a total surface area of 1.6 million km² at 20°-20° N longitude and 44°-64° E latitude. The political borders of Iran comprise a limited part of the orogenic belt (Zagros, Alborz and other mountain chains) that span the Arabian-African unit and the Asian block (Berberian & King 1981).

- Geomorphology and geology

The Zagros is a section of the Alpine-Himalayan orogenic belt and is the most extensive mountain range system in Iran and has a NW-SE orientation that stretches about 2000 km from eastern Turkey to the Makran Mountains. It provides a geomorphological separation between the Iranian plateau to the northeast and the Mesopotamian and Persian Gulf basins to the southwest. The system has a mean elevation of ~1305 m

with the highest point being 4408 m at Zard Kooh peak in Bakhtiari province (Homke 2007). The Alborz mountain range is an active, arcuate fold-and-thrust belt (Stocklin 1974, 1977, Axen *et al.* 2001). As the northern part of the Alpine-Himalayan belt, its gently sinuous east-west range extends into the north of Iran below the Caspian Sea. This system faces the depressed Caspian block to the north and is confined to the central Iranian plateau to the south (Stöcklin 1974). This mountain chain is about 950 km in length and its width varies from 15 to 110 km (Ghorbani 2013). The average elevation of >2000 m has as its highest peak Mt. Damavand, a dormant volcano with 5670 m in elevation.

Another important orogenic belt is Kopet-Dagh that stretches from the eastern margins of the Caspian Sea into northeastern Iran, Turkmenistan, and northern Afghanistan (Afshar-Harb 1979, Buryakovsky *et al.*

2001). Other mountains ranges in Iran include the Makran in central, Jebal Barez in Southeastern, and Azarbaijan in northwestern Iran. Interior basins, lowlands, coastal, and riverine zones are surrounded by these mountain systems (Fischer 1968).

Iran's habitat is classified as upland with an average altitude of 1000 m, with the exception of the limited plains of the Caspian Sea, Persian Gulf, and Khuzestan (Ghorbani 2013). Elevation varies from 27 m below sea level in the Caspian basin to 5671 m above sea level at Mt. Damavand. Iran, as a section of the Alpine-Himalayan orogenic belt (Ghorbani 2013) and has been divided into 10 structural zones (Stoklin & Nabavi 1973). A series of more recent studies (Nabavi 1978, Alavi 1994) recognizes the structural zones of central Iran including Sanandaj-Sirjan, Zagros, Alborz, Azarbaijan, eastern Iran, southeastern Iran, and Kopet-Dagh (Fig. 1).




Fig. 1. a. Simple geological map of Iran, b. Geomorphological map of Iran (www.ngdir.ir).

- Climatology

Climate is a chief component of ecology and affects the growth of plants and their distribution patterns. Other environmental factors, such as land-forms, habitat types, and soil characteristics demonstrate their effects locally (Zahran 2010). The topographic heterogeneity of Iranian ecosystems exhibit great diversity of climatic conditions. Mountainous massifs enclose the central parts of the country as a natural barrier and prevent penetration of moisture-bearing clouds into the interior of central Kavir and Dasht-e Loot. These orographic features experience variation in rainfall

(Djamali *et al.* 2011) as followed: Abali (2465 m asl, northeastern Iran, 534 mm), Astara (18 m bsl, northwestern Iran, 1380 mm), Bandar Abbas (10 m asl, southern Iran, 182.5 mm), Chabahar (8 m asl, southeastern Iran, 111 mm), Tabas (711 m asl, central Iran, 83 mm), Ilam (1337 m asl, western Iran, 616 mm) and Mashhad (999 m asl, northeastern Iran, 255.2 mm).

Djamali *et al.* (2011) classified Iranian ecosystems into macro-bioclimatic as follows: temperate (north), Mediterranean (west, northwest), and tropical (south accordant to zones of the Persian Gulf and Gulf of Oman). Breckle (2002) classifies Iranian habitats as

situated in zonobiom III to the south and zonobiom VII (rIII) to the north. The climatic heterogeneity, floral history, evolutionary potential (Frey & Probst 1986), complex interaction between climatic zones, and biogeographical regions in the orographic context (Zohary 1973) has produced many heterogenic habitats and plant formations (Frey & Probst 1986). Flora of Iran comprise 7300 plant taxa (Akhani 2006) and 1727 endemic species (Jalili & Jamzad 1999), and it has made Iran as an important center of endemism (Heywood 1994) with valuable zones of plant diversity (Barthlott *et al.* 1996).

Iran is an arid land with an annual rainfall that is less than one-third of the world average (Shakur *et al.* 2010). The hot southern gulf region, with its higher winter and summer temperatures and scant rainfall, displays a climate regimen similar to that of the tropical northeast African and the hot Sindian deserts, but with occasional more extreme maxima and minima (Zohary 1973).

Materials and Methods

The present study examined 3123 herbarium specimen records from HSBU, W and WU (herbarium abbreviation according to Thiers 2008) and several scientific resources (Schönbeck-Temesy 1972, Khatamsaz 2002) on the vegetation and flora of Iranian habitats to provide distribution and ecological data on *Solanaceae* in Iran. Climate factors were provided by the Iranian Climatological Organization (average of 50 years (1955–2005) for rainfall and temperature. Rivas-Martínez *et al.* (1997, 1999) developed a novel method of global classification for detailed bioclimatic analysis. Djamali *et al.* (2011) adapted it for the Iranian ecosystems that used in this study. Assessment of the conservation status was based on the IUCN Red List at regional scale (IUCN 2011). We classified Iranian plant formations based on Frey & Probst (1976) and plant habitats based on a combination of Zohary (1973) and Bailey (2009).

An ecogeographical database was developed using these records that corresponds to 2560 localities. The localities were marked using ArcView version 3.2

(ESRI 2000) on geo-referenced maps ($1/10^6$) of Iran. The distribution patterns of the taxa were mapped per $1^\circ \times 1^\circ$ universal transverse Mercator grid cells (100 km² with the exception of boundary area) to determine the distribution pattern and the most important areas for conservation management.

The criteria for scoring were the index of species rarity in the study area (RI) based on Williams *et al.* (1996), index of species distribution (SDE) based on Selvi (1997), Sapir *et al.* (2003), and Solymos & Feher (2005). Scoring ranged from zero (0) to one (1) for both indices and the vulnerability of the species increases as the score increases. The RI is calculated as the inverse of the number of cells having species recorded the target area as $RI=1/C_i$, where C_i is the number of grid cells and l is the number of present taxa. The SDE is calculated as $SDE=l-C_i/C$ where C is the total number of grid cells. To calculate the conservation value (CV), the RI and n (SDE of each grid cell) were summed up. The higher scores represent a higher CV. Biological forms were assessed based on Raunkaier (1934). Flora Iranica (Schönbeck-Temesy 1972) was the most important reference used for assessment and taxonomical nomenclature.

This is worthy to state that, all taxonomical information given here is merely on conservation-based approach, as the basic needs of the survey carried out in the present study.

Results

- Diversity and distribution patterns

Solanaceae is widely distributed in Iran. The family includes 55 species belonging to nine genera classified into six tribes (Schönbeck-Temesy 1972, Khatamsaz 2002, Akhani & Ghorbani 2003, Dinarvand & Howeizeh 2104) (Table 1). The taxa occur in diverse abundance in Iranian habitats, especially in the mountainous habitats of Taftan (East), central Alborz (North), and western slopes of central Zagros (West). Jebal Barez shows the greatest abundance and central Kavir shows the least at the family level (Figs 3–4).

The *Solanum* taxa are crop wild relatives (up to 13 per grid cell) in the western zones of the central Alborz,

central Zagros, and Taftan (Figs 3–4), *Atropa* (up to 11 per grid cell) in central Alborz, *Datura* (up to 8 per grid cell) in central Alborz and western zones of central Zagros (Fig. 2), *Withania* (up to 13 per grid cell) in Makran and central Taftan (Figs 3–4), *Lycium* (up to 19 per grid cell) in Kopet Dagh, northern and southern Zagros, and Taftan (Figs 3–4), and *Hyoscyamus* (up to 41 per grid cell). Central Alborz and Kopet Dagh show the highest frequency. *Solanaceae* (up to 55 per grid cell) shows the highest abundance in central Alborz, Kopet Dagh and Taftan. The highest richness of species and genera among the geographical latitudes are 14–15 (35°–37°) and 6 (36°–37°), respectively. The lowest number of species and genera per grid cell is one and reflects an irregular pattern in Iran. The species and genera vary from 1 to 15 and 1 to 6 per grid cell, respectively. The highest richness of species and genera is focused in the Alborz and Zagros regions.

This family is distributed over 155 of 171 grid cells as follows: *Hyocyanus* (110), *Lycium* (109), *Solanum* (71), *Withania* (25) *Atropa* (7), *Datura* (32), *Mandragora* (2), *Physochlaina* (Schönbeck-Temesy 1972) (1) (Figs 3–4). The genera distribution by geographical latitude is: *Hyocyanus* 26°–40°, *Lycium*

26°–39°, *Solanum* 26°–39°, *Withania* 26°–32°, *Atropa* 36°–38°, *Datura* 27°–40° and *Mandragora* 37°–38° and 32°–33° (Fig. 2). Moreover, the genera and species show irregular variation over the latitudinal gradient. A maximum of 1–15 species and a minimum of 1–5 genera occupy each grid cell. These taxa are diversely abundant in Iranian habitats. Taftan (East), central Alborz (north), the western slopes of central Zagros (west) and Jebal Barez show the greatest abundance. The central Kavir represents the lowest abundance at the family level (Figs 3–4). *Solanum* (up to 13 in grid cells) is found in western zones of the central Alborz, central Zagros and Taftan (Figs 3–4), *Atropa* (up to 11 in grid cells) in the central Alborz, *Datura* (up to eight in grid cells) in central Alborz and western zones of central Zagros (Figs 3–4), *Withania* (up to 13 grid cells) in Makran and central Taftan (Figs 3–4), *Lycium* (up to 19 in grid cells) in Kopet Dagh, the northern and southern Zagros and Taftan (Figs 3–4), *Hyocyanus* (up to 41 in grid cells) in the central Alborz and Kopet Dagh showed the greatest abundance. *Solanaceae* (up to 55 in grid cells) showed the greatest abundance in central Alborz, Kopet Dagh and Taftan.




Fig. 2. a. Species profile along latitude, b. Genera profile along latitude.




Fig. 3. Abundance pattern of the genera of *Solanaceae* in scale of Iran: a. *Hyocymus*, b. *Solanum*, c. *Atropa*, d. *Datura*, e. *Lycium*, f. *Withania*, g. *Solanaceae*.




Fig. 4. a. Richness of species, b. Richness of genera inside occupied grid cell.

The elevation patterns of the taxa can be classified as alpine (>2500 m), sub-alpine (1200–2500 m) and mountain slopes-to-lowlands (<1200 m), (Kapos *et al.* 2000), although many taxa show transitional habitats and occupy all three life zones. The taxa represent a range of

elevations as follows: *Solanaceae* (*Solanum* species only) 6–3650 m, *Physaleae* 10–1800 m (comprising *Physalis* at 10–1750 m and *Withania* at 10–1800 m), *Atropeae* 0–2200 m (comprising *Lycium* at 0–2200 m, *Atropa* at 8–1960 m, and *Mandragora* at 185 m)

Datureae at 1400 m, *Hyoscyamine* at 5–3800 m, *Physochlaina* at 640 m (Schönbeck-Temesy 1972), *Hyoscyamus* at 10–3800 m (Fig. 6). *Solanaceae* is distributed across an altitudinal range of 0 to 3800 m within the Iranian geographic borders.

Solanaceae shows distribution patterns in quaternary deposits and sedimentary, volcano-sedimentary, metamorphic, igneous, and ophiolitic rock. *Atropa* occurs in sedimentary and volcano-sedimentary rock only (simplified geological map of Iran www.ngdir.ir). The other genera inhabit a wide range of distribution in geological rock. *Solanaceae* is distributed in diverse geomorphological units in northern, northwestern, northeastern and central Iran, the Zagros, the coastlines of the Persian Gulf and Caspian Sea, and southern, southwestern and southeastern Iran. *Withania* is located in southwestern and southern geomorphological units and the others are established in a varied scope of geomorphological units.

The life forms of the plant taxa illustrate adaptation to environmental agents (Lande 1982, Galan de Mera *et al.* 1999). These forms represent the peak of adaptation of a plant to climatic conditions (Raunkiare 1908), therefore, assessment and analysis the life forms can help identify ecological features of the habitats (Diaz & Cabido 1997). *Solanaceae* in Iran features therophytes with 12 (19%), hemicryptophytes with 28 (45.7%), and chamaephytes with 25 (35.3%) (Fig. 7). *Solanum* features the highest abundance rates of hemicryptophytes (8), chamaephytes (4) and therophytes (4). The life forms diversity showing a peak at 1000–1500 m and sharply decrease over 2000 m. Besides, showing a peak between 29° to 32° latitude and with a clear decrease at the higher and lower latitudes (Fig. 7).

- Endemism

Most endemic taxa are belong to *Hyoscyamus* including *Hyoscyamine kotschyanus*, *H. malekianus*, *H. orthocarpus*, *H. natans*, and *H. rosularis* at 27° to 38° N latitude and 15 to 3800 m in elevation. This covers the Irano-Turanian and Sudano-Zambeian regions accordant to the Mediterranean pluviseasonal-continental, Mediterranean desertic-continental, and

tropical desertic bioclimates. *H. kotschyanus* is found in metamorphic and sedimentary rock, *H. malekianus*, *H. orthocarpus*, and *H. rosularis* in sedimentary rock and quaternary deposits, *H. natans* in sedimentary and igneous rock and infrequently in metamorphic rock.

- Rare taxa and conservation status

Rare taxa show greater extinction risk than common taxa in similar conditions of ecology (Johnson 1998, Matthies *et al.* 2004) because of high sensitivity to demographic and environmental events (Boyce 1992). *Solanaceae* taxa were, categorized based on RI. The rare classes were the very rare (10) to rare (7) and comprise 37% of Iranian *Solanaceae*. These taxa are distributed across the different bioclimatic, geomorphologic, and geologic units of Iran. The species occur at 18 to 3800 m in elevation. The rare taxa are found in *Solanum*, *Atropa*, *Mandragora*, *Physochlaina*, and *Hyoscyamus* genera.

Assessment of the conservation status of the plant taxa and developments of red lists are the first step in management. The IUCN Red List of threatened species is the most respected base for assessment of conservation categories. Endemic species are distributed from 8 to 3800 m.

- Phytogeography

Solanaceae include several widespread, cosmopolitan, and some bi-regional taxa. This family is found in the Irano-Turanian region (Holarctic kingdom) in the northern, western, eastern, and central parts and the Sudano-Zambeian region (Pleotropic Kingdom) in the southern parts. The first region comprises the Irano-Turanian regional center of endemism and second region comprises the Saharo-Sindian local center of endemism (Léonard 1989, White and Léonard 1991). The Irano-Turanian region shows clear climatic, vegetative, and floristic properties characterized by diurnal and annual variation in temperatures, low precipitation, hot summers, and cold extreme winters. This region shows little affinity to neighboring regions that featured with climax vegetation dominated by hemicryptophytes and chamaephytes (Zohary 1973) The Irano-Turanian (48.07%) and multi-regional elements (32.69%) have the highest contribution of phytochorions. The

Hyrceanian (7.69%), Irano-Turanian-Hyrceanian (5.76%), cosmopolitan (2%), and Sudano-Zambezian (1.88%) comprise other phytogeographical elements.

- Priorities for conservation

Myers (1988, 1990) classified hotspots for regions with specific concentrations of species richness, restricted endemics, and threatening factors. The factors considered for selection as priorities for conservation follow:

a. Richness-based

Solanaceae shows a regular pattern of species richness in the central Alborz, Zagros, Kopet Dagh and Makran mountains in Iran and have the highest rank for CV.

b. Rarity-based

The taxa classified that ranked highest for rarity have the highest priority for conservation. The most highly ranked are *Solanum* (6), *Hyoscyamus* (3), *Atropa* (2), *Mandrogora* (2), and *Physochlaina* (1), respectively.

c. Endemism-based

Endemism- taxa of *Solanaceae* all belong to *Solanum* (1), and *Hyoscyamus* (7) that are mainly centered in the Sudano-Zambezian region.

d. Habitat-specificity based

This category refers to the habitat for each taxon based on SDE. The taxa can be classified as high specificity, medium specificity, low specificity, and widespread.

- Conservation value

CV ranged from 0.15 (coast of Persian Gulf) to 11.19 (central Alborz) show an irregular pattern in Iran (Fig. 8).


Discussion

This is the first study on the distribution patterns of *Solanaceae* in Iran based on the geographic information system (GIS). The family comprises 55 species (0.82% of Iranian plant taxa) with eight endemic

species (0.46% of Iranian endemic taxa) (Hedge & Wendelbo 1978, Jalili & Jamzad 1999). This is a higher rate of endemism than in neighboring countries in southwest Asia. The center of origin of *Solanaceae* in South America (Hunziker 1979) shows a wide distribution (Barroso *et al.* 1991) and several species are distributed in limited and specific habitats around the world (Albuquerque *et al.* 2006).

Some insist on a Gondwanan origin for *Solanaceae* (e.g. Symon 1991) and others (Raven & Axelrod 1974) believe it is even younger. The high power dispersal and variance are important reasons for modern-day distributions (D'Arcy 1991). The prominent zones of endemism for *Solanaceae* are located in the Zagros mountainous ecosystem, the center of endemism of the Irano-Turanian region (Davis *et al.* 1994, Jalili & Jamzad 1999, Mehrabian *et al.* 2015), and the ecotonic zones (Clements 1905, Risser 1995) at the crossroads of the Irano-Turanian and Sudano-Zambezian regions.

The Irano-Turanian and Mediterranean regions (Takhtajan 1986, Zohary 1973) show higher diversity and endemism than adjacent areas (Clement, Hochstrasser & Peters). These centers encompass the richness of endemic monocotyledons in Iran (Mehrabian *et al.* 2015). *Hyoscyamus* L. and *Solanum* L. are endemic at 15 to 3800 m and create a wide and sensitive conservative zone along this elevation gradient. Geology is highly effective for delimitation of ecosystems and analysis of distribution patterns (Parenti & Ebach 2009). The endemic zones of *Solanaceae* are situated mainly in sedimentary (quaternary deposits) and ophiolitic formations in Iran. Geological evidence confirms the belief of Rechinger (1961) that specific soils can be used to delineate high endemism in Iran. Endemic *Solanaceae* shows a slight decrease from the Iranian plateau (Schönbeck-Temesy 1972) to the Anatolian plateau (Davis 1978) (Fig. 5).

Fig. 5. Endemism pattern of *Solanaceae* in SW Asia.Table 1. A comparison of *Solanaceae* members among some country of SW Asia

Genus	Iran	Iraq	Afghanistan	Pakistan	Russia	Syria and Palestine	Turkey	Saudi Arabia
<i>Solanum</i>	14	4	8	12	22	6	8	1
<i>Physalis</i>	2	-	1	1	6	1	1	-
<i>Hyoscyamus</i>	19	11	7	4	8	5	7	1
<i>Atropa</i>	4	-	1	1	3	1	1	-
<i>Physochlaina</i>	1	-	-	-	3	-	1	-
<i>Lycium</i>	7	3	4	6	5	2	8	2
<i>Datura</i>	2	1	3	3	3	2	3	2
<i>Mandragora</i>	2	-	-	-	1	1	1	-
<i>Withania</i>	2	1	2	2	-	1	1	1
Total	54	20	26	29	52	19	31	7

Table 2. Conservation and rarity of studied taxa. SDI (Index of Species Distribution), RI (Rarity Index) and CV (Conservation Value)

Taxon	SDE	RI	CV	Taxon	SDE	RI	CV
<i>Solanum pseudocapsicum</i>	0.984	1	1.984	<i>Atropa acuminata</i>	0.938	0.142	1.081
<i>S. nigrum</i>	0.476	0.029	0.506	<i>Mandragora turmanica</i>	0.892	1	1.892
<i>S. carmanicum</i>	0.984	1	1.984	<i>M. autumnalis</i>	0.892	1	1.892
<i>S. transcasicum</i>	0.984	1	1.984	<i>Datura stramonium</i>	0.984	0.043	1.028
<i>S. olgae</i>	0.907	0.166	1.074	<i>D. innoxia</i>	0.646	0.066	0.712
<i>S. alatum</i>	0.846	0.1	0.946	<i>Physochlaena orientalis</i>	0.984	1	1.984
<i>S. luteum</i>	0.876	0.125	1.001	<i>Hyoscyamus arachnoideus</i>	0.415	0.026	0.441
<i>S. kieseritzkii</i>	0.953	0.333	1.287	<i>H. turcomanicus</i>	0.753	0.062	0.816
<i>S. persicum</i>	0.600	0.038	0.638	<i>H. niger</i>	0.553	0.034	0.588
<i>S. asiae-mediae</i>	0.969	0.5	1.469	<i>H. kurdicus</i>	0.661	0.045	0.706
<i>S. incanum</i>	0.984	0.06	1.051	<i>H. leucanthera</i>	0.784	0.071	0.856
<i>S. dulcamara</i>	0.769	0.047	0.816	<i>H. squarrosus</i>	0.676	0.047	0.724
<i>S. sisymbriifolium</i>	0.907	0.5	1.407	<i>H. reticulatus</i>	0.476	0.029	0.506
<i>S. surattense</i>	0.953	0.2	1.153	<i>H. pojarkovae</i>	0.969	0.5	1.469
<i>Physalis alkekengi</i>	0.923	0.090	1.013	<i>H. kotschyanus</i>	0.938	0.25	1.188
<i>P. divaricate</i>	0.830	0.125	0.955	<i>H. pusillus</i>	0.138	0.017	0.156
<i>Withania somnifera</i>	0.876	0.090	0.967	<i>H. senesionis</i>	0.461	0.028	0.490
<i>W. coagulans</i>	0.830	0.047	0.878	<i>H. leptocalyx</i>	0.984	1	1.984
<i>Lycium ruthenicum</i>	0.676	0.027	0.704	<i>H. malekianus</i>	0.969	0.5	1.469
<i>L. depressum</i>	0.446	0.015	0.461	<i>H. rosularis</i>	0.984	1	1.984
<i>L. kopetdaghi</i>	0	0.076	0.076	<i>H. tenuicaulis</i>	0.676	0.047	0.724
<i>L. dasystemum</i>	0.8	1	1.8	<i>H. orthocarpus</i>	0.830	0.090	0.921
<i>L. makranicum</i>	0.984	0.2	1.184	<i>H. insanus</i>	0.584	0.037	0.621
<i>L. shawii</i>	0.923	0.033	0.956	<i>H. nutans</i>	0.846	0.1	0.946
<i>L. edgeworthii</i>	0.538	0.062	0.600	<i>H. bonniillerianus</i>	0.984	1	1.984
<i>Atropa belladonna</i>	0.753	0.2	0.953				
<i>A. pallidiflora</i>	0.923	1	1.923				
<i>A. komaroxii</i>	0.984	0.25	1.234				




Fig. 6. Genera richness in altitudinal profile: a. *Solanum*, b. *Atropa*, c. *Lycium*, d. *Physalis*, e. *Hyocymus*, f. *Datura*.




Fig. 7. Comparative life form diversity in: Altitudinal profile (a), Latitudinal profile (b), Wholly richness variation of *Solanaceae* along latitudinal (c) and altitudinal (d) profiles. (Th: Therophyte, Ch: Chamaephyte, Hem: Hemicryptophyte).

The genus *Lycium* L. comprises about 70 species in a disjointed distribution in arid and semi-arid regions and the saline habitats of coastlines in temperate and subtropic zones in America, Africa, and Eurasia (Hitchcock 1932, D’Arcy 1979, 1991, Hunziker 1979). *Lycium* in Iran (7) and Eurasia (10) shows less diversity

than in North (20) and South America (30) and North Africa (20). The disjointed dispersal of *Solanaceae* has been hypothesized as being caused by continental drift (Symol 1991) or long-distance dispersal (Raven & Axelord 1974). *Lycium* forms known plant communities in Irano-Turanian (*L. kopetdaghi*, *L. ruthenicum*), and

Sudano-Zambezian (*L. shawii*, *L. makranicum*) regions in Iran (Zohary 1973). Some are umbrella taxa (Groom *et al.* 2006, Christopher 2007) that support lower plant communities. They are influenced by wildlife that feed upon their fruit and shape their refuge (Mostafavi *et al.* 2006, Abdoli *et al.* 2008) to conserve biodiversity. They are surrogates of biodiversity (Santi *et al.* 2010, Chiarucci *et al.* 2015) and keystones to sustainability and protect the ecosystem (Paine 1995, Mills *et al.* 1993). Zohary (1973) classified *Lycium shawi* as an important element in steppe-scrub of *Juniperus-Pistacia-Amygdalus* and oligo-hydric communities along the coastlines of the Persian Gulf and Sea of Oman. Modern species are considered indicators of former connections between Iran and east Africa through tropical Arabia (Zohary 1973) as the red fruit is easily dispersed by birds over long distances. *Lycium kopetdaghi*, and *L. ruthenicum* are Irano-Turanian elements and, in their eastern-most distribution range, cover the northeast and east, respectively.

Atropa is a forest-sub-forest element distributed in limited localities of Hyrcanian forests. *Atropa belladonna* is situated in the most eastern portions of the distribution range and *A. acuminata* in the most western portion of the distribution range in Iran. Unfortunately several populations of recent taxa have failed during the widespread destruction of the Hyrcanian forests.

Hyoscyamus, comprising about 20 species with a distribution throughout Eurasia, except for disjunctions caused by deserts (Tu *et al.* 2010). This genus is represented by 19 species (as a center of diversity) in Iran, including seven endemic species. Tu *et al.* (2005) have been suggested that divergence among *Hyoscyameae* may have resulted from the rise of the Himalayan-Hengduan mountain ecosystem alongside the Tethys margins and the extension the dryness in central Asia. Polyploidy is a common occurrence in *Hyocymaeae* and plays a role in harsh habitats (Hanelt 1966). It is also common in *Hyoscyameae*, especially in the *Hyocymus* that restricted to Iran (Sheidai *et al.* 1998). Some have classified them into the following anthropogenic communities in Iran (Zohary 1973, Khatamsaz 1998):

H. reticulatus, *H. Turcomanicus*, and *H. arachnoides*. The negative effects of agricultural ecosystems means that distribution data on recent taxa is necessary to their conservation and management (Mack *et al.* 2000).

Mandragora L. is an old world taxa of *Solanaceae* that include 6 species (Dinarwand & Howizeh 2014) distributed in the circum-Mediterranean, Irano-Turanian, and Sino-Himalayan regions (Ungriht *et al.* 1998). *M. turcomanica* Mizg is a neglected and underutilized plant (IPGRI 2002) with a limited distribution in Turkmenistan (Kurbanov 1994) from a small zone in Golestan in northern Iran (Akhani & Ghorbani 2003), and *M. autumnalis* Bercol have recently been reported by Dinarwand & Howizeh (2014) in oak forests in the Shimbar Preserve in Khuzestan (SW Iran). Traditional societies of Iran could be the reason for the decrease in distribution because of overharvesting. Establishment of a preserve has given *M. autumnalis* a better chance of conservation. *M. turcomanicum* is found in home gardens (Watson *et al.* 2001) and the support of local offices of the environment can improve protection by the owners. Penetration of this Mediterranean element into western regions of Iran confirms the belief of Zahran (2010) and Zohary (1973) that the western slopes of the Zagros have been affected by the Mediterranean region. A sister affinity between two recent taxa as well as their disjunction patterns (in the eastern Mediterranean basin and Irano-Turanian region) empowered a hypothesis that includes vicariance of Mediterranean flora and fauna (Kadereit & Yapark 2008). The Mediterranean-Himalayan dispersal of *Mandragora* may have been caused by periodical fluctuations in the Caspian Sea during the Pliocene and Quaternary that simulated divergence and allopatric speciation in recent taxa as well the following taxa: *Mespilus germanica*, *Sorbus graeca*, *Evonimus velutina*, *Crataegus androssowi*, and *Rhus coriaria* (Fet & Atamuradov 1994). Others (Proskurayakova & Belyanian 1985, Kurbanov 1994) insist on the role of fauna on their dispersal and believe on a younger distribution. In traditional societies of Iran, they are called "Mehr-e giah", which could be the reason for the overharvest and decrease in the distribution range

in the region. Regardless of its limited distribution in Iran, establishment of a protected area has preserved the conservation status of *M. autumnalis*.

The genus *Datura* includes 10 to 12 species (Geeta and Gharaibeh 2007) that are mainly distributed in southern North America (Bye 1986). *D. ferox* L., *D. stramonium*, and *D. metel* L. are distributed in China, Eurasia and south and west Asia, respectively. Symon (1991) insisted that the origin of diversification *Datura* is in the New World.

Solanum is the largest genera in *Solanaceae* and one of the largest taxa of angiosperms (Olmstead *et al.* 1999) and shows less diversity in Iran than in other regions of its distribution. Several *Solanum* species are distributed mainly in Iranian protected areas (Fig. 9). These taxa are located in highly enriched zones that can be suggested as gene sanctuaries (Singh 1981) or gene micro-reserves (Serra *et al.* 2004) for protection of these valuable wild relatives of crops. They are valuable for breeding plants for resistance to disease and environmental stresses and potential uses for agronomy (Jansky 2000). *Solanum* subgenus *Leptostemonum* (Dunal) Bitter. is a large group of monophylytes (ca. 350–450 taxa) with worldwide dispersal centered dominantly in South and Central America, Africa, Australia and Asia. It can be recognized by the presence of prominent epidermal thorns that cover the stems and leaves (Levin *et al.* 2006, Olmstead 2013). *Solanum incanum* is a member of this subgenus and is a common species in association with *Ziziphus spina-christii* in the Sudano-Zambian in the Middle East, especially in Iran (Zohary 1973). This stretches to Saudi Arabia, Africa, Pakistan and India. Another spiny species (*Solanum surattense*) distributed in southeastern Iran shows wide distribution in America, Europe, Africa and southwest Asia (Levin *et al.* 2006). *Solanum sisimbryfolium* Lam. (spiny *Solanum*) is a newly penetrated taxon recorded from the lowlands of the Caspian Sea and is widely distributed in ruderal habitats along roadsides and sand dunes in northern Iran (Eslami & Naqinezhad 2010 and personal observation of second author). This recent species is native to South America but has naturalized in

Europe, some habitats of Africa and Australia as well as having a wide distribution in the US (Karaer & Kutbay 2007). This pioneer taxon was dispersed through the fleshy and red fruit by birds (Hill & Hulley 2000) and mice (Bryson 2011). *Solanum sisimbryfolium* appears to have invasive effects in Africa (Hill & Hulley 1995, Byrne *et al.* 2002), and US (USDA 2013). The ability of this species to spread to new areas indicates that, ecological considerations should be considered to control it in relict Hyrcanian forests.

Physalis with about 70 species is centered in Mexico and is endemic to this region. Only two of these species occur jointly in America and Asia (*P. divaricata*) along with an isolated Asiatic-European species (*P. alkegeni*) (Olmstead *et al.* 2008) distributed in northern, western, and southern parts of Iran. Substantial changes in natural habitats mean these rare taxa are faced with a reduction in distribution range and population size to the point where only a few individuals have been observed in reported sites. Overharvesting for medicinal use is also a threat in Iran.

Withania has 10–18 species (based on diverse systematic classifications) and is distributed throughout the Canary Islands, the Mediterranean region and northern Africa to India, China, and Japan (Hepper 1991, Hunziker 2001). Some species, especially *W. somnifera* have ethno-botanical uses and medicinal properties (Chevallier 1996, Darias *et al.* 2001). Modern species have been classified as farm grown (segetal) in tropical areas in the Middle East (Zohary 1973).

Iran is located at the crossroads of diverse phytochorions (Zohary 1973). A wide spectrum of phytogeographical elements applies to the *Solanaceae* of Iran. Irano-Turanian (48.07) and multi-regional elements (32.69) contribute most significantly to phytochorions. The Hyrcanian (7.69), Irano-Turanian-Hyrcanian (5.76), cosmopolitan (2), and Sudano-Zambeian (1.88) are other phytogeographical elements. The Mediterranean elements distributed in the northern forests of Iran (Hyrcanian) confirm recent studies (Rechinger 1989, Akhiani & Ziegler 2002).

The 21 taxa (40.7%) of *Solanum* (8), *Hyoscyamus* (7), *Lycium* (2), *Atropa* (3), and *Physalis*, each represented by only one species and are the most threatened (critically endangered and endangered) taxa of *Solanaceae* in Iran. A high percentage of threatened taxa are situated in preserves. The central Alborz, central Zagros, and some central mountains of Iran cover the most important conservative zones for *Solanaceae* in Iran. The most threatening factors of land-use change, deforestation, and overgrazing place serious pressure on the habitats of these taxa. Providing a conservation plan for protection of the endangered taxa is necessary. *Ex situ* methods can be used for conservation of several taxa of *Solanaceae* (Sponer *et al.* 2000, Sarasan *et al.* 2006) and are important support for improvement of crops against biotic and abiotic stress.

The present study established the most important centers of diversity for *Solanaceae* in Iran accordant to central Asia and the Middle East (Vavilov 1951). Fortunately, a broad area is situated inside Iranian Protected Areas (Bali & Bahmapour 2012). *Solanaceae* in Iran shows greater diversity of species/genera (53/9) than Afghanistan (26/7), Pakistan (29/7), Syria, Palestine (19/8), and Turkey (36/12) in southwestern Asia, with the exception of the USSR (14:62). The species *Solanum* and *Hyoscyamus* are clearly declining from eastern and western parts of southeastern Asia (Afghanistan and Pakistan) to eastern parts of southwest Asia (Syria and Palestine). Ecotonical conditions caused by penetration of the Mediterranean climate (Zahran 2010) and the intersection of the Irano-Turanian and Mediterranean phytochorions (Takhtajan 1986) is a possible reason behind its richness in the Zagros. Alborz and the Majoran mountains are centers of Iranian plant diversity (Davis *et al.* 1994) and an important part of the Saharo-Sindian local center of endemism (Zohary 1973). The Mediterranean and tropical bioclimatic units (Djamali *et al.* 2011) are optimal climates for *Solanaceae* as in Europe and South America.

Analysis shows that the richness of *Solanaceae* undergoes minor changes at 1500 m and sharply decreases at above 1500 m. *Solanum* and *Hyoscyamus*

have the widest world distribution (Simpson 2007) that reaches the highest elevations and geographic range of distribution. *Nicaandrea* has a reported limited range (0–500 m) in Iran. At 0–500 m, 40 species form the highest and at 3500–4500 m, four species form lowest zones of richness for *Solanaceae*. It mainly occupies lowlands to semi-mountainous habitats in Iran, but shows wide distribution across the elevations. Their habitats are primarily igneous and sedimentary-volcanic sediment (Iranian Department of Geology).

Several taxa of *Solanaceae* crop wild relatives (e.g. *Solanum*, *Hyocyanus*, *Datura*, *Atropa*, and *Mandragora*) have valuable potential for improvement of the crop gene pool (Anjula *et al.* 2008). They should be clearly mapped for selection of important zones as possible plant micro-reserves (Heywood and Dullo 2005) and genetic reserves in Iran. These zones cover parts of two important centers of diversity in central Asia and the Middle East based on the Vavilov classification (Ladizinsky 1998).

Solanaceae classified in anthropogenic communities of Iran (Zohary 1973, Khatamsaz 1998) are *Hyoscyamus reticulatus*, *H. turcomanicus*, *H. archnoides*, *Datura inoxia*, and *D. stramonium*. Several taxa of *Solanaceae* as crop wild relatives have valuable potential for improvement of the crop gene pool (Anjula *et al.* 2008). The results of this study map important zones for the possible establishment of plant micro-reserves (Heywood & Dullo 2005), genetic reserves, and gene sanctuaries (Sing 1981, Laguna & Lumberas 2001, Tan & Tan 2002) for *Solanaceae* in Iran.

Life forms show adaptation of plant taxa to environmental agents (Lande 1982, Galan de Mera *et al.* 1999) and ecological features of habitats (Diaz & Cabido 1997). These forms characterize the peak of adaptation of a plant to climatic conditions (Raunkiare 1908) and as an adaptive strategy to hydrothermic gradients plays an important role in distribution patterns (Wang *et al.* 2002).

Hemicryptophytes at 46.7% and therophytes at 20% are the highest and lowest spectra of life forms for *Solanaceae*. Hemicryptophytes decrease sharply above

2000 m asl compared with chamaephytes and therophytes. Hemicyptophytes at 46.7% and therophytes at 20% cover the highest and lowest spectra of life forms in *Solanaceae*. Moreover, therophytes with highest richness between 1000–1500 m asl and at 29°–32° in latitude decrease sharply at above 2000 m asl (Fig. 8). The peak lies in the southern segment of the endemic zone (Hedge & Wendelbo 1978) as well southern zone of center of plant diversity (Boulus *et al.* 1994) in Iran.

Almost half of Iranian *Solanaceae* appear to have very rare to rare distributions and face several challenges to survival. Assessment of the conservation status of the plant taxa is a preliminary step in management plans and selection of priorities for conservation. Threatened categories of Iranian *Solanaceae*, including CR (43.4%) and VU (34%), mainly (58.9%) distributed in Irano-Turanian phytochorion. They are distributed at <2500 m asl, which is a sensitive zone that must be included in conservation plans. Severe destruction of habitat has been caused by overgrazing and land use changes (Mehrabian 2014) and *Solanaceae* is considered to be a threatened family on the regional scale of Iran. Disregard to this, only *M. turcomanicum* with a restricted

distribution in northeastern Iran and southern Turkmenistan is critically endangered on the global scale. Recent taxa in northeastern Iran are situated in home gardens only; therefore, urgent support, notification by the local offices of the environment, and establishment of micro-reserves seems necessary for *in situ* conservation.

Myers (1988, 1990) classified hotspots for regions with specific concentrations of species richness, restricted endemics, and threatened taxa. Rarity (Williams *et al.* 1996) and conservation value (Tsifits *et al.* 2009) are the other criteria for selection of priorities. When resources for conservation are narrow, identification priorities play a key role in conservation (Myers 2000, Brummitt & Lughadha 2003). In Iran, *Solanaceae* represents the highest richness in the central Alborz, Zagros, Kopet Dagh and Makran Mountains based on recent criteria. Based on endemism, the mountainous ecosystem of the central Zagros as the most important center of Irano-Turanian region should be included in these priorities. The conservation value of habitats ranges from 0.15 (coastline of Persian Gulf) to 11.19 (eastern Alborz) and show irregular patterns in Iran (Table 2).




Fig. 8. Conservation value of different parts of Iran based on distribution of *Solanaceae* species.




Fig. 9. Iranian Protected Areas (Bali & Bahmanpour 2012).

A considerable percentage of threatened taxa are situated in Iranian protected areas; however, several factors continue to be a threat to them, including land use change, deforestation and overgrazing. A conservation plan to protect recent taxa seems necessary. *Ex situ* methods have been used for conservation of several members of *Solanaceae* (Sponer *et al.* 2000, Sarasan *et al.* 2006) as complementary to *in situ* protection. These achievements create valuable data with which to manage the endemic, rare, crop-wild relatives and medicinal taxa

References

- Abdoli, A., Kiabi, B., Liaghati, H., Mostafavi, H., Mehrabian, A.R., Khoshbakht, K. & Mousavi, B. 2008. Atlas of Bushehr province Ecotourism. Shahid Beheshti University Publishing. 144 pp.
- Afshar, A. 1979. Stratigraphy, tectonic and petroleum geology of Kopet-Dagh region, northeast Iran: Doctoral College of Science and Technology. University of London, England, 316 pp.
- Agard, P., Omrani, J., Jolivet, L. & Mouthereau, F. 2005. Convergence history across Zagros, Iran; constraints from collisional and earlier deformation: International Journal of Earth Sciences 94: 401–419.
- Aghanabati, A. 1998. Jurassic Stratigraphy of Iran. Vol. 1. Geological Survey of Iran, Tehran.
- Akademische, D., Verlagsanstalt, G.A. & Symon, D.E. 1991. Gondwanan elements of the *Solanaceae*. Pp. 139–150. In: J.G. Hawkes, R.N. Lester, M. Nee & R. Estrada (eds). *Solanaceae* III. Taxonomy, Chemistry, Evolution. Royal Botanic Gardens, Kew.
- Akhani, H. 1998. Plant biodiversity of Golestan National Park, Iran. Stapfia 53: 1–411.
- Akhani, H. 2006. Flora Iranica: facts and figures and a list of publications by K.H. Rechinger on Iran and adjacent areas. Rostaniha 7(Suppl. 2): 19–61.

of *Solanaceae* and prioritize habitats and geographical zones to design conservation plans for this family in Iran.

Acknowledgments

This paper is part of a larger study entitled “Diversity, distribution patterns, conservation status and priorities for conservation to select important plant areas in Iran” that was funded by the Research Council of Shahid Beheshti University, Tehran, Iran (Grant No. 600/1829, 600/3226, 600/3513). We like to thank Mr. Ali Moazeni for his help with the GIS maps.

- Akhani, H. & Ziegler, H. 2002. Photosynthetic pathways and habitats of grasses in Golestan National Park (NE Iran) with an emphasis on the C4-grass dominated rock communities. *Phytocoenologia* 32(3): 455–501.
- Akhani, H. & Ghorbani, A.B. 2003. *Mandragora turcomanica* (*Solanaceae*) in Iran: a new distribution record for an endangered species. *Systematics and Biodiversity* 1(2): 177–180.
- Alavi, N.M. 1994. Tectonic of the Zagros, orogenic belt of Iran, new data and interpretation. *Tectonophysics* 299: 211–238.
- Albuquerque, L.B., Velázquez, A. & Mayorga-Saucedo, R. 2006. *Solanaceae* composition, pollination and seed dispersal syndromes in Mexican Mountain Cloud Forest: *Acta Botanica Brasilica* 20(3): 599–613.
- Alejandra Jaramillo, M. & Paul, S. 2001. Phylogeny and patterns of floral diversity in the genus *Piper* (*Piperaceae*). *American Journal of Botany* 88(4): 706–716.
- Anderson, S. 1994. Area and endemism. *Quarterly Review of Biology* 69: 451–471.
- Andrew, R., Marshall Philip, J., Platts, R., Gereau, E., Kindeketa, W., Kang'ethe, S. & Marchant, R. 2012. The genus *Acacia* (*Fabaceae*) in East Africa: Distribution, diversity and the protected area network. *Plant Ecology and Evolution* 145(3): 289–301.
- Anjula, P., Ajay Kumar Tomer, Bhandari D.C., Pareek S.K. 2008. Genetic Resources and Crop Evolution. Towards collection of wild relatives of crop plants in India. 55: 187–202.
- Applequist, W., Mcglinn, D.J., Miller, M., Long, Q.G. & Miller, J.S. 2007. How well do herbarium data predict the location of present populations? A test using *Echinacea* species in Missouri. *Biodiversity and Conservation* 16: 1397–1407.
- Arponen, A., Heikkinen, R., Thomas, C. & Moilanen, A. 2005. The value of biodiversity in reserve selection: representation, species weighting, and benefit functions. *Conservation Biology* 19(6): 2009–2014.
- Atamuradov, K. & Fet, V. 1994. Biogeography and Ecology of Turkmenistan (Monographiae Biologicae, Vol. 72), Natural Conservation Society, Ashgabat, Turkmenistan, Department of Biological Sciences, Loyola University, New Orleans, Louisiana, USA.
- Axen, G.J., Lam, P.S., Grove, M., Stockli, D.F. & Hassanzadeh, J. 2001. Exhumation of the west-central Alborz Mountains, Iran, Caspian subsidence and collision-related tectonics. *Geology* 29: 559–562.
- Bali, A. & Bahmanpoor, A. 2012. The Distribution Map of Iranian Protected Areas. Department of Environment.
- Ball, P.W. 1972. *Onosma* L. "Flora Europea. Vol. 3. *Diapensiaceae* to *Myoporaceae*", pp. 89–94.
- Barroso, G.M., Peixoto, A.L., Ichaso, C.L.F., Costa, C.G., Guimarães, E.F. & Lima, H.C. 1991. Sistemática de Angiospermas do Brasil. v. 3. Viçosa, Universidade Federal de Viçosa.
- Barthlott, W., Lauer, W. & Placke, A. 1996. Global distribution of species diversity in vascular plants: Towards a world map of phytodiversity. *Erdkunde* 50: 317–328.
- Barthlott, W., Mutke, J., Braun, G. & Kier, G. 2000. Die ungleiche globale Verteilung pflanzlicher Artenvielfalt-Ursachen und Konsequenzen. *Berichte der Reinhold Tüxen. Gesellschaft* 12: 67–84.
- Barthlott, W., Hostert, A., Kier, G., Kuper, W., Kreft, H., Mutke, J., Rafiqpoor, M.D. & Sommer, J.H. 2007. Geographic patterns of vascular plant diversity at continental to global scales. *Erdkunde* 61(4): 305–316.
- Berberian, M. 1983. The southern Caspian: A compressional depression floored by a trapped, modified oceanic crust: *Canadian Journal of Earth Sciences* 20: 163–183.
- Berberian, M. & King, G.C.P. 1981. Towards a paleogeography and tectonics evolution of Iran.

- Canadian Journal of Earth Sciences 18(2): 210–265.
- Bernau, S. & Heider, S. 2015. *Ex situ* Conservation Priorities for the Wild Relatives of Potato (*Solanum* L. Section *Petota*). PLOS ONE 10(6): e0129873.
- Bibby, C.J., Collar, N.J., Crosby, M.J., Heath, M.F., Imboden, C., Johnson, T.H., Long, A.J., Stattersfield, A.J. & Thirgood, S.L. 1992. Putting Biodiversity on the Map: Priority Areas for a Global Conservation. ICBP, Cambridge.
- Bischler, H. & Jovet-Ast, S. 1986. The hepatic flora of South-West Asia: a survey. Proceedings of the Royal Society of Edinburgh. Section B. Biological Sciences 89: 229–241.
- Boissier, E. 1879. Flora Orientalis sive enumeratio plantarum in Oriente a Graecia et Aegypto and Indiae fines hucusque observatarum. IV: Coroliflorae et Monochlamydeae. Geneva: H. Gorge.
- Braun, G., Mutke, J., Reder, A. & Barthlott, W. 2002. Biotope patterns, phytodiversity and forestline in the Andes, Based on GIS and Remote Sensing Data. In: C. Körner & E.M. Spehn (eds). Mountain Biodiversity: a Global Assessment. Parthenon Publishing, London. Pp. 75–89.
- Breckle, S.W. 2002. Walter's Vegetation of the Earth. The Ecological Systems of the Geo-biosphere. Springer, Heidelberg, 527 pp.
- Brummitt, N. & Lughadha, E.N. 2003. Biodiversity: where is hot and where is not hot? Conservation Biology 17: 1442–1448.
- Buryakovsky, L.A., Chilingir, G.V. & Aminzadeh, F. 2001. Petroleum geology of the South Caspian Basin: Gulf Professional Publishing USA, 442 pp.
- Byrne, M.J., Currin, S. & Hill, M.P. 2002. The influence of climate on the establishment and success of the biocontrol agent *Gratiana spadiccea*, released on *Solanum sisymbriifolium* in South Africa. Biological Control 24: 128–134.
- Caro, T. 2010. Conservation by Proxy. Island Press, USA.
- Cecchi, L. & Selvi, F. 2009. Phylogenetic relationships of the monotypic genera *Halascya* and *Paramoltkia* and the origins of serpentine adaptation in *Circum mediterranean Lithospermeae* (*Boraginaceae*): insights from ITS and matK DNA sequences. Taxon 58: 700–714.
- Chiarucci, A., D'Auria, F., De Dominicis, V., Lagan'a, A., Perini, C. & Salerini, E. 2005. Using vascular plants as a surrogate taxon to maximize fungal species richness in reserve design. Conservation Biology 1644–1652.
- Clements, F.E. 1905. Research methods in ecology. University Publishing Company, Lincoln, NE.
- D'Arcy, W.G. 1979. Classification of *Solanaceae*. In: The Biology and Taxonomy of the *Solanaceae* (G.J. Hawkes, R.N. Lester & A.D. Skelding, eds). Pp. 3–47. Academic Press, London.
- D'Arcy, W.G. 1991. The *Solanaceae* since 1976 with a review of its biogeography. In: *Solanaceae* III: Taxonomy-Chemistry-Evolution (J.G. Hawks, R.N. Lester, M. Nee & N. Eserada, eds). Royal Botanic Garden, Kew and the Linnean Society of London, Richmond. Pp. 75–137.
- Davidson, J., Hassanzadeh, J., Berzins, R., Stockli, D.F., Bashukoo, B., Turrin, B. & Pandamouz, A. 2004. The Geology of Damavand Volcano, Alborz Mountains, northern Iran. Geological Society of America Bulletin 116(1–2): 16–29.
- Davis, P.H. 1978. Flora of Turkey 6. Edinburgh: Edinburgh University Press.
- Davis, S.D., Heywood, V.H. & Hamilton, A.C. 1994. Centres of Plant Diversity. A Guide and Strategy for their Conservation. Vol. 1. Europe, Africa, South West Asia and the Middle East. IUCN and WWF, Gland, Switzerland. 354 pp.
- Davis, S., Heywood, V.H. & Hamilton, A.C. 1994–97. Worldwide Fund for Nature and International Union for Conservation of Nature and Natural Resources, Centers of Plant Diversity (Gland, Switzerland).
- Delisle, F., Lavoie, C., Jean, M. & Lachance, D. 2003. Reconstructing the spread of invasive plants:

- taking into account biases associated with herbarium specimens. *Journal of Biogeography* 30: 1033–1042.
- Dinarvand, M. & Howeizeh, H. 2014. A new record of the genus *Mandragora* (*Solanaceae*) for the flora of Iran. *The Iranian Journal of botany* 20(2): 179–182.
- Djamali, M., Akhiani, H., Khoshravesh, R., Anderieu-Ponel, P. & Brewer, S. 2011. Application of the global bioclimatic classification to Iran: implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea* 37(1): 91–114.
- Dobson, A.P., Rodriguez, J.P., Roberts, W.M. & Wilcove, D.S. 1998. Geographic Distribution of Endangered Species in the United States. Reid, W. V. Biodiversity hotspots. *Trends in Ecology & Evolution* 13: 275–280.
- Droissart, V., Hardy, J.O., Sonke, B., Dahdouh Guebas, F. & Ste´vart, T. 2012. Subsampling Herbarium Collections to Assess Geographic Diversity Gradients: A Case Study with Endemic *Orchidaceae* and *Rubiaceae* in Cameroon. *Biotropica* 44(1): 44–52.
- Eftekharnajad, J. 1980. Tectonic classification of Iran based on the pattern of sedimentary basins. *Bulletin of Petrological Society* 82: 19–28 (In Persian)
- Eslami, B. & Naqinezhad, A. 2010. A new shrubby *Solanum* (*Solanaceae*) species for the flora of Iran. *The Iranian Journal of Botany* 17(2): 254–256.
- ESRI. 2000. ArcView Gis Ver. 3.2a. Environmental Systems Research Institute Inc., California.
- Faith, D.P. & Walker, P.A. 1996. Environmental diversity: on the best-possible use of surrogate data for assessing the relative biodiversity of sets of areas. *Biodiversity and Conservation* 5: 399–415.
- Farjon, A. 1998. World checklist and biogeography of conifers. Royal Botanical Garden, Kew.
- Fisher, W.B. 1968. Physical Geography. In: W.B. Fisher (ed.). *The Cambridge History of Iran*, Cambridge University Press 1: 3–110.
- Franklin, J. 2009. Mapping species distributions: spatial inference and prediction. Cambridge University Press, Cambridge.
- Fuentes, N., Sa´nchez, P., Esquivel, J. & Marticorena, A. 2012. A new comprehensive database of alien plant species in Chile based on herbarium records. *Biological Invasions*. Pp. 847–858.
- Gaston, K.J. 1996. Species richness: Measure and measurement. *Biodiversity: A biology of numbers and difference*, ed. K.J. Gaston. Cambridge, MA: Blackwell Science. Pp. 77–113.
- Ghorbani, M. 2013. *The Economic Geology of Iran*. Springer. 581 pp.
- Groom, M.J., Meffe, G.K. & Carroll, C.R. 2006. *Principles of Conservation Biology*. Third edition. Sinauer Associated, Sunderland.
- Hansen, A.J. & DiCasteri, F. 1992. Landscape boundaries: consequences for biotic diversity and ecological flows. *Ecological Studies* 92. Springer-Verlag, New York.
- Hawkes, J.G., Lester, R.N., Nee, M. & Estrada-Ramos, N. 1992. *Solanaceae* III. Taxonomy, Chemistry, Evolution. Royal Botanic Gardens, Kew. Pp. 785–788.
- Hawkins, B.A., Field, R. & Cornell, H.V. 2003. Energy, water, and broad scale geographic patterns of species richness. *Ecology* 84: 3105–3117.
- Hedge, C. & Wendelbo, P. 1978. Patterns of Distributions and Endemism in Iran. *Notes from the Royal Botanical Garden* 36: 441–464.
- Herna´ndez, H.M. & Navarro, M. 2007. A new method to estimate areas of occupancy using herbarium data. *Biodiversity and Conservation* 16: 2457–2470.
- Hernandez-Stefanoni, J.L. 2006. The role of landscape patterns of habitat types on plant species diversity of a tropical forest in Mexico. *Biodiversity and Conservation* 15: 1441–1457.
- Heiser, C.B. 1969. *Nightshades, the Paradoxical Plants*. San Francisco: Freeman & Company.
- Hill, M.P. & Hulley, P.E. 1995. Biology and host range of *Gratiana spadicea* (Klug 1829) (*Coleoptera: Chrysomelidae: Cassidinae*), a potential

- biological control agent for the weed *Solanum sisymbriifolium* Lamarck (*Solanaceae*) in South Africa. *Biological Control* 5(3): 345–352.
- Hill, M.P. & Hulley, P.E. 2000. Aspects of the phenology and ecology of the South American weed, *Solanum sisymbriifolium*, in the Eastern Cape province of South Africa. *African Plant Protection* 6(2): 53–59.
- Hilty, J.A., Lidicker, W.A. & Merenlender, M. 2006. *Corridor Ecology: The Science and Practice of Linking Landscapes for Biodiversity Conservation*. Island Press, Washington.
- Hitchcock, C.L. 1932. A monographic study of the genus *Lycium* of the western hemisphere. *Annals of Missouri Botanical Garden* 19: 179–374.
- Heywood, V.H. & Dulloo, M.E. 2005. *In situ* conservation of wild plant species: a critical global review of best practices. IPGRI Technical Bulletin 11. IPGRI, Rome, Italy.
- Hobohm, C. 2003. Characterization and ranking of biodiversity hotspots: centers of species richness and endemism. *Biodiversity and Conservation* 12: 279–287.
- Houshmand Zadeh, A. 1998. Geology of the Basement of Central Iran.
- Hunziker, A.T. 1979. South American *Solanaceae*: A synoptic survey. *In: The Biology and Taxonomy of the Solanaceae* (G.J. Hawkes, R.N. Lester & A.D. Skelding, eds). Academic Press, London. Pp. 49–85.
- IUCN. 2011. Guidelines for appropriate uses of IUCN Red List Data. Incorporating the Guidelines for Reporting on Proportion Threatened and the Guidelines on Scientific Collecting of Threatened Species. Version 2. Adopted by the IUCN Red List Committee and IUCN SSC Steering Committee. 78 pp.
- Iriondo, J.M., Maxtend, N. & Dulloo, M.E. 2008. *Conserving Plant Genetic Diversity in Protected Areas*. CAB International Publication, London, UK.
- IUCN Standards and Petitions Subcommittee. 2011. *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 9.0. Prepared by the Standards and Petitions Subcommittee.
- Jalili, A. & Jamzad, Z. 1999. *The Red Data Book of Iran*. Research Institute of Forests and Rangelands, 748, Tehran.
- Janick, J. 2002. *History of Horticulture*: Purdue University. <http://www.hort.purdue.edu/newcrop/history/lecture30/lec30.html>.
- Jansky, S.H. 2000. Breeding for disease resistant in potato. *Plant Breeding Review* 19: 69–155.
- Jedicke, E. 2001. Biodiversität, Geodiversität, Ökodiversität. Kriterien zur Analyse der Landschaftsstruktur-ein konzeptioneller Diskussionsbeitrag. *Naturschutz und Landschaftsplanung* 33: 59–68.
- Jetz, W., Rahbek, C. & Colwell, R.K. 2004. The coincidence of rarity and richness and the potential signature of history in centres of endemism. *Ecology Letters* 7: 1180–1191.
- Kadereit, G. & Yaprak, A.E. 2008. *Microcnemum coralloides* (*Chenopodiaceae-Salicornioideae*): an example of intraspecific East-West disjunctions in the Mediterranean region. *Anales del Jardín Botánico de Madrid* 65: 415–426.
- Kala, C.P. 2005. Indigenous Uses, Population Density, and Conservation of Threatened Medicinal Plants in Protected Areas of the Indian Himalayas: *Conservation Biology* 19: 368–378.
- Kandari, L.S., Rao, K.S., Maikhuri, K., Kharkwal, G., Chauhan, K. & Kala, C.P. 2011. Distribution pattern and conservation of threatened medicinal and aromatic plants of Central Himalaya, India: *Journal of Forestry Research* 22(3): 403.
- Kapos, V., Rhind, J., Edwards, M., Price, M.F., Ravilious, C. & Butt, N. 2000. Developing a map of the world's mountain forests. Forests in sustainable mountain development: a state of knowledge report for 2000. Task Force on Forests in Sustainable Mountain Development, pp. 4–19.

- Karaer, F. & Kutbay, H.G. 2007. *Solanum sisymbriifolium* Lam. (*Solanaceae*): a new record for Turkey. Turkish Journal of Botany 31: 481–483.
- Kati, V., Devillers, P., Dufre ne, M., Legakis, A., Vokou, D. & Lebrun, P. 2004. Hotspots, complementarity or representativeness? Designing optimal small-scale reserves for biodiversity conservation. Biology Conservation 120: 471–480.
- Khatamsaz, M. 2002. *Boraginaceae* L. In: Assadi *et al.* (eds). Flora of Iran. Vol. 39, pp. 114–168. Research Institute of Forests and Rangelands, Iran (In Persian).
- Kolarc ik, V. & Ma rtonfi, P. 2006. Revision of the distribution of the genus *Onosma* (*Boraginaceae*) in Slovakia. Thaiszia. Thaiszia 16: 131–154.
- Kolarc ik, V., Zozomova Lihova, V.J.J. & Ma rtonfi, P. 2010. Systematics and evolutionary history of the *Asterotricha* group of the genus *Onosma* (*Boraginaceae*) in central and southern Europe inferred from AFLP and nrDNA ITS data. Plant Systematic Evolution 290: 21–45.
- Kreft, H. & Jetz, J. 2007. Global patterns and determinants of vascular plant diversity. PNAS 104(14): 5925–5930.
- Kurbanov, D. 1994. Flora of Kopetdagh. In: V. Fet & K.I. Atamuradov (eds). Biogeography and Ecology of Turkmenistan. Kluwer Academic Publishers, Dordrecht, pp. 105–128.
- Kuznetzow, N.I. 1913. *Boraginaceae*. In: N.I. Kuznetzow, N.A. Bush & A.V. Fomin (eds). Flora Caucasica Critica, Vol. 4, No. 2, K. Mattisena, Jurijev, pp. 66–400.
- Lavoie, C. 2013. Biological collections in an ever changing world: Herbaria as tools for biogeographical and environmental studies. Perspectives in Plant Ecology, Evolution and Systematics 15: 68–76.
- Ladiges, P.Y., Marks, C.E. & Nelson, G. 2011. Biogeography of *Nicotiana* section *Suaveolentes* (*Solanaceae*) reveals geographical tracks in arid Australia, Journal of Biogeography 38: 2066–2077.
- Ladizinsky, G. 1998. Plant Evolution under Domestication. The Netherlands: Kluwer Academic Publishers. Blaine P. Friedlander Jr. (2000-06-20). Retrieved 2008-03-19.
- Leslie Dowe, J. 2010. Biogeography, Ecology and Systematic Csiro. Scientific and Industrial Research Organisation.
- L eonard, J. 1981–89.   l etude de la flore et de la v eg tation des d serts d’Iran:  tude des aires de distribution les phytochories, les chorotypes. Fascicules 1–9. Meise, Jardin botanique national de Belgique.
- Levin, R.A., Myers, N.R. & Bohs, L. 2006. Phylogenetic relationships among the “Spiny Solanums” (*Solanum* subgenus *Leptostemonum*, *Solanaceae*). American Journal of Botany 93(1): 157–169.
- Linder, H.P. 2001. Plant diversity and endemism in sub-Saharan tropical Africa. Journal of Biogeography 28: 169–182.
- Lovett, J.R., Rudd, S., Taplin, J. & Frimodt-Moller, C. 2000. Patterns of plant diversity in Africa south of the Sahara and their implications for conservation management. Biodiversity and Conservation 9: 37–46.
- Mack, R.N., Simberloff, D., Lonsdale, W.M., Evans, H., Clout, M. & Bazzaz, F.A. 2000. Biotic invasions: causes, epidemiology, global consequences and control. Ecological Society of America 10: 689–710.
- McNaughton, S.J. 1994. Conservation Goals and the Configuration of Biodiversity. In: P.L. Forey, C.J. Humphries & R.I. Vane-Wright (eds). Systematics and Conservation Evaluation. The Systematic Association (Special issue), Vol. 50. Clarendon Press, Oxford.
- Meusel, H., J ager, E., Rauschert, S. & Weinert, E. 1978. Vergleichend Chronologie der zentrieuropalschen Flora-Kareten, Band II. Gustav Fischer Verlag. Jena.
- Mehrabian A.R., Sheidai, M., Noormohammad, Z., Mozafarian, V. & Asri, Y. 2012. Interpopulations diversity in *Onosma microcarpa* (*Boraginaceae*):

- Morphological and molecular approach. *Science Medical* 3: 187–198.
- Michael, J. & Hopkins, G. 2007. Modeling the known and unknown plant: biodiversity of the Amazon Basin. *Journal of Biogeography* 34: 1400–1411.
- Mittermeier, R.A., Myers, N., Thomsen, J.B. & da Fonseca, G.A.B. 1998. Biodiversity hotspots and major tropical wilderness areas: approaches to setting conservation priorities. *Conservation Biology* 12: 516–520.
- Mittermeier, R.A., Myers, N., Gil, P.R. & Mittermeier, C.G. 1999. Hotspots: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions (*CEMEX*, Conservation International and Agrupacion Sierra Madre, Monterrey, Mexico).
- Mittermeier, R.A., Gil, P.R., Hoffman, M., Pilgrim, J., Brooks, T., Mittermeier, C.G., Lamoreux, J. & da Fonseca, G.A.B. 2005. Hotspots revisited: Earth's biologically richest and most endangered ecoregions. Conservation International (CI), Washington, DC.
- Moreira-Muñoz, A. & Muñoz-Schick, M. 2007. Classification, diversity, and distribution of Chilean *Asteraceae*: implications for biogeography and conservation. *Diversity and Distributions* 13: 818–828.
- Mostafavi, H., Kiabi, B., Abdoli, A., Mehrabian, A.R., Mahini, A.R., Kami, H. & Naqinezhad, A.R. 2006. Biodiversity of Mond Protected Area. Shahid Beheshti University Publishing, 254 pp.
- Mozaffarian, V. 1378. Flora Khuzestan: Markaz-e Tahghighat-e Manabe Tabiee va Dam-e Khuzestan, 670 pp. (In Persian).
- Mozaffarian, V. 1379. Flora of Yazd: Moasese-ye Entesharat-e Yazd, 636 pp. (In Persian).
- Müller, R., Nowicki, C., Barthlott, W. & Ibich, P.L. 2003. Biodiversity and endemism mapping as a tool for regional conservation planning-case study of the *Pleurothallidinae* (*Orchidaceae*) of the Andean rain forests in Bolivia. *Biodiversity and Conservation* 12: 2005–2024.
- Mutke, J. & Barthlott, W. 2005. Patterns of vascular plant diversity at continental to global scales. *In*: I. Friis & H. Balslev (eds). Plant diversity and complexity patterns-Local, Regional and Global Dimensions. The Royal Danish Academy of Sciences and Letters, Copenhagen, Biologiske Skrifter 55: 521–537.
- Myers, N., Mittermeier, R.A., Mittermeier, C.G. & da Fonseca, G.A.B. 2000. Biodiversity hotspots for conservation priorities, *Nature* 403.
- Nabavi, M. 1978. An introduction to geology of Iran. Geological Survey of Iran, Tehran.
- Negi, H.R. & Gadgil, M. 2002. Cross-taxon surrogacy of biodiversity in the Indian Garhwal Himalaya. *Biological Conservation* 105: 143–155.
- Niels, R., Marco, C., Roos, J.W., Ferry, S.E. & Loon, E. 2009. Botanical richness and endemic patterns of Borneo derived from species distribution models. *Ecography* 32: 180–192.
- Nogole-Sadat, M.A.A. 1978. Les zones de décrochement et les virgation structurales en Iran. Consequences des resultants de l'analyse structural de la région du Qom. Unpublished PhD thesis, France, 201 p.
- Noroozi, J., Akhiani, H. & Breckles, S.W. 2008. Biodiversity and Phytogeography of Alpine Flora of Iran. *Biodiversity and Conservation* 17(2): 493–521.
- Olmstead, R.G. 2013. Phylogeny and biogeography in *Solanaceae*, *Verbenaceae* and *Bignoniaceae*: a comparison of continental and intercontinental diversification patterns. *Botanical Journal of Linnean Society* 171: 80–102.
- Olmstead, R.G., Sweere, J.A., Spangler, R.E., Bohs, L. & Palmer, J.D. 1999. Phylogeny and provisional classification of the *Solanaceae* based on chloroplast DNA. Pp. 111–137. *In*: M. Nee, D. Symon, R.N. Lester & J. Jessop (eds). *Solanaceae IV: Advances in Biology and Utilization*. Royal Botanic Gardens, Kew.
- Ortega-Baes, P. & Godínez-Alvarez, H.C. 2006. Global diversity and conservation priorities in the

- Cactaceae*: Biodiversity and Conservation 15: 817–827.
- Olson, D.M. & Dinerstein, E. 1998. The Global 200: A representative approach to conserving the Earth's most biologically valuable ecoregions. *Conservation Biology* 12: 502–515.
- Parenti, L.R. & Ebach, M.C. 2009. *Comparative Biogeography*. University of California Press.
- Panaia, M., Senaratna, T., Bunn, E., Dixon, K.W. & Sivasithamparam, K. 2000. Micropropagation of the critically endangered Western Australian species, *Symonanthus bancroftii* (F. Muell.) L. Haegi (*Solanaceae*): *Plant Cell Tissue and Organ Culture* 63: 23–29.
- Pedrotti, F. 2013. *Plant and Vegetation Mapping*. Springer, Heidelberg.
- Peralta, I.E. & Spooner, D. 2007. History, origin and early cultivation of tomato (*Solanaceae*). Genetic improvement of Solanaceous crops. M.K. Razdan & A.K. Mattoo. Enfield (NH), Science Publisher 2: 1–24.
- Petrus, C. & Cunninghamia, H. 1998. Some New South Wales coastal plant distributions: a comparison of herbarium records with transect survey data. *Cunninghamia* 5(3): 645–664.
- Prendergast, J.R., Quinn, R.M., Lawton, J.H., Eversham, B.C. & Gibbons, D.W. 1993. Rare species, the coincidence of diversity hotspots and conservation strategies. *Nature* 365: 335–337.
- Priscilla, H.C. & Bruce, W. 2009. Can herbarium records be used to map alien species invasion and native species expansion over the past 100 years? *Journal of Biogeography* 36: 651–661.
- Rabinowitz, D. 1981. *Biological Aspects of Rare Plant Conservation*. Wiley, New York, pp. 205–217.
- Raunkaier, C. 1905. Types biologiques pour la géographie botanique. *Oversigt over Det Forhandlinger*: 347–438.
- Raunkaier, C. 1934. *The life forms of plants and statistical plant geography*. Clarendon Press, Oxford, 632 pp.
- Raven, P.H. & Axelrod, D.I. 1974. Angiosperm biogeography and past continental movements. *Annals of Missouri Botany Garden* 61: 39–637.
- Rechinger, K.H. 1986. *Cousinia*: Morphology, taxonomy, distribution and phytogeographical implications. *Proceeding of the Royal Society of Edinburgh* 89B: 45–58.
- Rechinger, K.H. 1989. Fifty years of botanical research in the Flora Iranica Area (1937–1987). In: K. Tan (ed.). *Plant taxonomy, phytogeography and related subjects*. The Davis & Hedge Festschrift, Edinburgh, pp. 301–349.
- Regens, N. 1997. Floristic biodiversity and history of African arid regions *Biodiversity and Conservation* 6: 495–514.
- Reveal, J.L. 1981. The concept of rarity and population threats in plant communities. *Rare plant conservation*, ed. L.E. Morse & M.S. Henefin. Bronx: New York Botanical Garden. Pp. 41–46.
- Richard, G. & Olmstead, R.G. 2013. Phylogeny and biogeography in *Solanaceae*, *Verbenaceae* and *Bignoniaceae*: a comparison of continental and intercontinental diversification patterns. *Botanical Journal of the Linnean Society* 171: 80–102.
- Riedl, H. 1968. *Boraginaceae*. In: K.H. Rechinger (ed.). *Flora Iranica* 48. Akademische Druck-u. Verlagsanstalt. Graz: Austria.
- Risser, P.G. 1995. The status of the science of examining ecotones. *Bioscience* 45(5): 318–325.
- Rivas-Martínez, S., Sánchez-Mata, D. & Costa, M. 1997. Syntaxonomical synopsis of the potential natural plant communities of North America I. *Itinera Geobotanica* 10: 5–148.
- Rivas-Martínez, S., Sánchez-Mata, D. & Costa, M. 1999. Boreal and western temperate forest vegetation (syntaxonomical synopsis of the potential natural plant communities of North America II). *Itinera Geobotanica* 12: 3–311.
- Robenson, E. 2008. *Medicinal Plants at Risk. Nature's Pharmacy, Our Treasure Chest: Why We Must Conserve Our Natural Heritage*. A Native Plant

- Conservation Campaign Report. Center for Biological Diversity. 16 pp.
- Sapir, Y., Shmida, A. & Fragman, O. 2003. Constructing red numbers for setting conservation priorities of endangered plant species: Israeli flora as a test case. *Nature Conservation* 11: 91–107.
- Sarasan, V., Cripps, R., Ramsay, M.M., Atherton, C., McMichen, M., Prendergast, G. & Rowntree, J.K. 2006. Conservation *in vitro* of threatened plants-progress in the past decade. *In Vitro Cellular Developmental Biology* 42(3): 206–214.
- Sarfatti, G. 2005. Society for Conservation Biology. Dipartimento di Scienze Ambientali, Università degli Studi di Siena, Via P.A. Mattioli 4, 53100 Siena, Italy.
- Sankaran, M. 2009. Diversity patterns in Savanna grassland communities: implications for conservation strategies in a biodiversity hotspot. *Biodiversity Conservation* 18: 1099–1115.
- Santi, E., Maccherini, S., Rocchini, D., Bonini, I., Brunialti, G., Favilli, L., Perini, C., Pezzo, F., Piazzini, S., Rota, E., Salerni, E., Chiarucci, A. & Sarfatti, G. 2010. Simple to sample: Vascular plants as surrogate group in a nature reserve. *Journal for Nature Conservation* 18: 2–11.
- Scheiner, S.M. & Rey-Benayas, J.M. 1994. Global patterns of plant biodiversity. *Evolutionary Ecology* 8: 331–347.
- Schery, R.W. 1951. *Plants for Man*. Prentice Hall, Adapted from Vavilov Englewood Cliffs, NJ, 1972.
- Schmidt, M., Kreft, H., Thiombiano, A. & Zizka, G. 2005. Herbarium collections and field data-based plant diversity maps for Burkina Faso. *Diversity and Distributions* 11: 509–516.
- Schonbeck-Temesy, E. 1972. *Solanaceae*. In: K.H. Rechinger (ed.). *Flora Iranica*, Vol. 100, Akademische Druck-u. Verlagsanstalt. Graz, Austria.
- Schulman, L., Toivonen, T. & Ruokolainen, K. 2007. Analysing botanical collecting effort in Amazonia and correcting for it in species range estimation. *Journal of Biogeography* 34(8):1388–1399.
- Selvi, F. 1997. Rare plants on mount Aiata, Italy: vulnerability to extinction on an ecological “Island”. *Biology Conservation* 81: 257–266.
- Serra, L., Oérez-Rovira, P., Deltoro, V.I., Faabregat, C., Laguna, E. & Pérez-Botella, J. 2004. Distribution, status and conservation of rare relict plant species in the Valencian community. *Bocconea* 16: 857–863.
- Shakur, A., Roshan, G.h., Najafe, R. & Kanei, A. 2010. Evaluating climatic potential for palm cultivation in Iran with emphasize on degree-day index, *African Journal of Agricultural Research* 13: 99–118.
- Sheidai, M., Mosallanejad, M. & Khatamsaz, M. 1999. Karyological studies in *Hyoscyamus* species of Iran. *Nordic Journal of Botany* 19: 369–373.
- Simpson, M.G. 2007. *Plant systematics*, Elsevier Academic Press. 603 pp.
- Singh, B. 1981. Establishment of First Gene Sanctuary in India for *Citrus* in Garo Hills. Concept Publishing, New Delhi, India.
- Sołymos, P. & Feheér, Z. 2005. Conservation prioritization based on distribution of land snails in Hungary. *Conservation Biology* 19(4): 1084–1094.
- Sorensen, L.L. 2004. Composition and diversity of the spider fauna in the canopy of a montane forest in Tanzania. *Biodiversity Conservation* 13: 437–452.
- Spooner, D.M. & Hijmans, R.J. 2001. Potato systematics and germplasm collecting. *American Journal of Potato Research* 78: 237–268.
- Spooner, D.M., Van Den Berg, R.G. & Miller, J.T. 2000. Species and series boundaries of *Solanum* ser. *Longipedicellata* (*Solanaceae*) and phenetically similar species in ser. *Demissa* and ser. *Tuberosa*: implications for a practical taxonomy of Section *Petota*. *American Journal of Botany* 88(1): 113–130.
- Stattersfield, A.J., Crosby, M.J., Long, A.J. & Wege, D.C. 1998. *Endemic Bird Areas of the World*. Birdlife International, Cambridge.
- Stéphane, H. 2007. Timing of shortening and uplift of the Pusht-e kuh Arc in the Zagros,

- a combined magnetostratigraphy and Fold-and-Thrust Belt (Iran) Apatite Thermochronology Analysis. Group of Dynamics of the Lithosphere (GDL), Institut de Ciències de la Terra "Jaume Almera", CSIC.
- Stocklin, J. 1974. Northern Iran: Alborz Mountains. In: A.M. Spencer (ed.). Mesozoic-Cenozoic orogenic belts data for orogenic studies; Alpine-Himalayan Orogens: Geological Society of London (special publication) 4: 212–234.
- Stocklin, J. 1977. Structural correlation of the Alpine ranges between Iran and central Asia: Société Géologique de France, Memoire hors série 8: 333–353.
- Schmidt, M., Kreft, H., Thiombiano, A. & Zizka, G. 2005. Herbarium collections and field data-based plant diversity maps for Burkina Faso Diversity and Distributions. *Diversity and Distribution* 11: 509–516.
- Symon, D., Haegi, L.A.R. 1991. *Datura* (*Solanaceae*) is a New World genus. In: J.G. Hawkes, R.N. Lester, M. Nee & N. Estrada (eds). *Solanaceae* III: taxonomy, chemistry and evolution. Royal Botanic Gardens Kew, Richmond, pp. 197–210.
- Takhtajan, A. 1986. Floristic regions of the world. University of California Press, pp. 544, California.
- Takin, M. 1972. Iranian geology and continental drift in the Middle East: *Nature* 235: 147–150.
- Tchalenko, J.S. 1975. Seismicity and structure of the Kopet Dagh (Iran, U.S.S.R.). *Pyramid* 278 (1275): 1–28. Royal Society, London.
- Tie-Yao, T., Hang, S., Zhi-Jian, G. & Ji-Pei, Y. 2005. Cytological studies on the Sino-Himalayan endemic *Anisodus* and four related genera from the tribe *Hyoscyameae* (*Solanaceae*) and their systematic and evolutionary implications. *Botanical Journal of the Linnean Society* 147: 457–468.
- Thiers, B.M. 2008. Continuously updated: Index Herbariorum.
- Tilman, D. 1999. The ecological consequences of changes in biodiversity: a search for general principles. *Ecology* 80: 1455–1474.
- Tsiftsis, S., Tsiripidis, I. & Karagainnakidou, V. 2009. Identifying areas of high importance for orchid conservation in East Macedonia (NE Greece). *Biodiversity and Conservation* 18: 1765–1780.
- Tu, T., Volis, S., Dillon, M.O., Sun, H. & Wen, J. 2010. Dispersals of *Hyoscyameae* and *Mandragorae* (*Solanaceae*) from New World to Eurasia in the early Miocene and their biogeographic diversification within Eurasia *Molecular Phylogenetic and Evolution* 57: 1226–1237.
- Ungrecht, S., Knapp, S. & Press, J.R. 1998. A revision of the genus *Mandragora* (*Solanaceae*). *Bulletin of the Natural History Museum, London: Botany* 28(1): 17–40.
- USDA. 2013. Weed Risk Assessment for *Solanum sisymbriifolium* Lam. (*Solanaceae*). Sticky nightshade *Solanum*.
- Vane-Wright, R.I., Humphries, C.I. & Williams, P.H. 1991. What to protect? Systematics and the agony of choice. *Biology Conservation* 55: 235–254.
- Vavilov, N.I. 1951. The Origin, Variation, Immunity and Breeding of Cultivated Plants Trans. from the Russian by K. Starr Chester. Waltham, Mass.: *Chronica Botanica*. New York: Stechert-Hafner, 364 pp.
- Vega, I.L., Ayala, O.N.N. & Conteras-Medina, R.L. 2004. Patterns of diversity, endemism and conservation: an example with Mexican species of *Ternstroemiaceae* Mirb. ex DC. (*Tricolpates: Ericales*). *Biodiversity and Conservation* 13: 2723–2739.
- Wagenitz, G. 1986. *Centaurea* in South-West Asia: Patterns of distribution and diversity. *Proceedings of the Royal Society of Edinburgh. Sect. B. Biological Sciences*, 89B: 11–21.
- Watson, J.W. 2001. Home gardens and *In situ* conservation of plant genetic resources in farming systems. *Proceedings of the Second International*

- Home Gardens. Workshop, Witzenhausen, Federal Republic of Germany.
- White, F. & Léonard, J. 1991. Phytogeographical Links between Africa and Southwest Asia. *Flora et Vegetatio Mundi* 9: 229–246.
- Willis, J.C. 1973. A dictionary of the flowering plants and ferns. Cambridge University Press, Cambridge.
- Williams, P.H., Humphries, C.J. & Vane-Wright, R.I. 1991. Measuring biodiversity: taxonomic relatedness for conservation priorities. *Australian Systematic Botany* 4: 665–679.
- Williams, P.H., Gaston, K.J. & Humphries, C.J. 1997. Mapping biodiversity value worldwide: combining higher-taxon richness from different groups. *Proceeding of the Royal Society. London B* 264: 141–148.
- Zahraieva, M., Prospero, J.M. & Monneveux, P. 2004. Ecological distribution and species diversity of *Aegilops* L. genus in Bulgaria: Kluwer Academic Publishers.
- Zahran, M.A. 2010. Climate-Vegetation: Afro-Asian Mediterranean and Red Sea Coastal Lands. Springer.
- Zhongqoiang, L.I., Dan, Y.U., Wen, X., Dong, W. & Manghui, T.U. 2006. Aquatic plants diversity in arid zones of Northwest China: patterns, threats and conservation. *Biodiversity and Conservation* 15: 3417–3444.
- Zohary, M. 1973. Geobotanical foundations of the Middle East. 2 Vols. Stuttgart. 739 pp.

ROSTANIHA